arXiv:math/0404040v1 [math.GR] 2 Apr 2004

Relatively hyperbolic groups: Intrinsic geometry, algebraic properties, and algorithmic problems

D. V. Osin

Contents

1 Introduction **3**
- 1.1 Preliminary remarks . . . 3
- 1.2 Main results . . . 6

2 Relative isoperimetric inequalities **11**
- 2.1 Relative presentations and length functions . . . 11
- 2.2 Geometry of van Kampen diagrams over relative presentations . 14
- 2.3 Relative Dehn functions . . . 22
- 2.4 Splitting Theorem for relatively finitely presented groups . . . 29
- 2.5 Isoperimetric functions of Cayley graphs. . . . 35

3 Geometry of finitely generated relatively hyperbolic groups **40**
- 3.1 Conventions and notation . . . 40
- 3.2 Properties of quasi–geodesics . . . 42
- 3.3 Geodesic triangles in Cayley graphs . . . 51
- 3.4 Symmetric geodesics . . . 58

4 Algebraic properties **69**
- 4.1 Elements of finite order . . . 69
- 4.2 Relatively quasi–convex subgroups . . . 72
- 4.3 Cyclic subgroups and translation numbers . . . 79

5 Algorithmic problems **88**
- 5.1 The word and membership problems . . . 88
- 5.2 The parabolicity problems . . . 90
- 5.3 Algorithmic problems for hyperbolic elements . . . 94

Open questions **100**

Appendix. Equivalent definitions of relative hyperbolicity **103**

Bibliography **108**

Chapter 1

Introduction

1.1 Preliminary remarks

Originally, the notion of a relatively hyperbolic group was proposed by Gromov [45] in order to generalize various examples of algebraic and geometric nature such as fundamental groups of finite–volume non–compact Riemannian manifolds of pinched negative curvature, geometrically finite Kleinian groups, word hyperbolic groups, small cancellation quotients of free products, etc. Gromov's idea has been elaborated by Bowditch in [13]. (An alternative approach was suggested by Farb [37].) In the present paper we obtain a characterization of relative hyperbolicity in terms of isoperimetric inequalities and adopt techniques based on van Kampen diagrams to the study of algebraic and algorithmic properties of relatively hyperbolic groups. This allows to establish a background for the subsequent paper [68], where we use relative hyperbolicity to prove embedding theorems for countable groups.

Since the words 'relatively hyperbolic group' seem to mean different things for different people, we briefly explain here our terminology. There are two different approaches to the definition of the relative hyperbolicity of a group G with respect to a collection of subgroups $\{H_1, \ldots, H_m\}$. The first one was suggested by Bowditch [13]. It is similar to the original Gromov's concept and characterizes relative hyperbolicity in terms of the dynamics of properly discontinuous isometric group actions on hyperbolic spaces. (For exact definitions we refer to the appendix).

In the paper [37], Farb formulated another definition in terms of the coset graphs. In the simplest case of a group G generated by a finite set S and one subgroup $H \leq G$ it can be stated as follows. G is *hyperbolic relative to* H if the graph $\widehat{\Gamma}(G, S)$ obtained from the Cayley graph $\Gamma(G, S)$ of G by contracting each of the cosets gH, $g \in G$, to a point is hyperbolic. In fact, the hyperbolicity of $\widehat{\Gamma}(G, S)$ is independent on the choice of the finite generating set S in G.

The two definitions were compared in [80], where Szczepański showed that if a group G is hyperbolic with respect to a collection of subgroups $\{H_1, \ldots, H_m\}$ in the sense of Bowditch, then G is hyperbolic with respect to $\{H_1, \ldots, H_m\}$ in the sense of Farb, but not conversely. However, in [37] Farb do not simply consider relatively hyperbolic groups. He introduces an additional (and central in his theory) condition, the so–called Bounded Coset Penetration property (or BCP, for brevity). It turns out that the notion of the relative hyperbolicity with BCP in the sense of Farb is equivalent to the notion of the relative hyperbolicity in the sense of Bowditch [13, 21, 31].

In order to define relative hyperbolicity of a group G with respect to a collection of subgroups $\{H_1, \ldots, H_m\}$, the approaches of Farb and Bowditch require G to be finitely generated as well as $H_1, \ldots, H_m$ (although the last assumption is rather technical). In the present paper we introduce a more general definition which is based on relative isoperimetric inequalities. This allows us to eliminate the assumption of the existence of finite generation sets of G and $H_1, \ldots, H_m$ as well as the assumption of the finiteness of the collection of subgroups. Such a generalization is important in certain applications and, in particular, allows to include the small cancellation theory over free products (see [56, Ch. V, Sec. 9], [61, Ch. 11]) within the framework of the theory of relatively hyperbolic groups. In case the group G is finitely generated, our notion of relative hyperbolicity is equivalent to that of Bowditch and to that of Farb with the BCP property.

Before stating the main theorems, let us survey certain motivating examples for which, in particular, the results of our paper can be applied.

(I) Let M^n be a noncompact, complete, finite-volume Riemannian manifold with (pinched) negative sectional curvature

$$-b^2 \leq K(M^n) \leq -a^2 < 0.$$

Then $\pi_1(M^n)$ is hyperbolic in the sense of Bowditch with respect to the collection of cusp subgroups [37] (for the definition of cusp subgroups we refer to [34]). The examples of such a type include, for instance, non–uniform lattices in real $\mathbb{R}$–rank one simple Lie groups.

(II) Let G be a $C'(1/6)$–small cancellation quotient of the free product of groups $X_1, \ldots, X_k$ (see [56, Ch. V, Sec. 9]). Then G is hyperbolic relative to the natural images of subgroups X_i in G in the sense of Bowditch. This result follows directly from our characterization of relative hyperbolicity in terms of relative isoperimetric inequality and the Greendlinger Lemma [56] (see also [70]–[72], where 'hyperbolic factorizations' of free products are studied).

(III) Suppose that $H_1, H_2, \ldots, H_k$ are quasi–convex subgroups of a word hyperbolic group G. Then G is hyperbolic with respect to the collection $H_1, H_2, \ldots, H_k$ in the sense of Farb [42]. If, in addition, $|H_i^g \cap H_j| < \infty$ whenever $g \notin H_i$ or $i \neq j$, then G is hyperbolic relative to $H_1, H_2, \ldots, H_k$ in the sense of Bowditch [13, Theorem 7.11].

(IV) Yaman [84] characterized relatively hyperbolic groups as geometrically finite convergence groups (see Appendix for definitions). More precisely, suppose that M is a non–empty perfect compact metric space and suppose that a group G acts as a geometrically finite convergence group on M. Suppose also that the stabilizer of each bounded parabolic point is finitely generated. Then G is hyperbolic relative to the set of the maximal parabolic subgroups in the sense of Bowditch and M is equivariantly homeomorphic to the boundary of G. In the case of ordinary hyperbolic groups (i.e. in case the set of bounded parabolic points of M is empty) this result was obtained by Bowditch [11].

(V) Recently Dahmani [32] proved combination theorem for relatively hyperbolic groups. This allowed to show that the limit groups introduced by Sela [76] in his solution of the Tarski problem are hyperbolic in the sense of Bowditch relative to their maximal abelian non–cyclic subgroups (see also [2]).

We also mention some examples of groups, which are hyperbolic in the sense of Farb.

(VI) Let $Mod(S)$ denote the mapping class group corresponding to a surface S. Using the isometric action of $Mod(S)$ on the complex of curves introduced in [48], Masur and Minsky proved that $Mod(S)$ is hyperbolic relative to a finite collection of stabilizers of certain curves in the sense of Farb [59]. However, in most cases it is not hyperbolic in the sense of Bowditch. An alternative proof can be found in [14]

(VII) Applying a technique related to small cancellation theory, Kapovich [52] proved the relative hyperbolicity in the sense of Farb of some Artin groups of extra large type with respect to certain families of parabolic subgroups. Another result of this type was obtained by Bahls [4]. He showed that right-angled Coxeter groups are relatively hyperbolic in the sense of Farb with respect to natural collections of rank 2 parabolic subgroups.

(VIII) Finally we mention two combination theorems. If G is an HNN–extension of a group H (respectively an amalgamated product of H_1 and H_2) with associated (respectively amalgamated) subgroups A and B, then G is hyperbolic in the sense of Farb relative to H (respectively relative to $\{H_1, H_2\}$. Moreover, if H is hyperbolic in the sense of Farb relative to $\{A, B\}$ (respectively H_1 is hyperbolic in the sense of Farb relative to A and H_2 is hyperbolic in the sense of Farb relative to B), then G is hyperbolic in the sense of Farb relative to A [67]. In particular, this allows to construct a finitely presented group G which is hyperbolic in the sense of Farb relative to a hyperbolic subgroup H and has undecidable word problem.

1.2 Main results

In this section we discuss shortly the main results of our paper. We assume the reader to be familiar with such notions as Cayley graph, Dehn function, hyperbolic group, quasi–geodesic path, etc., and refer to the next chapter for the precise definitions.

Let G be a group, X a subset of G, and H an arbitrary subgroup of G. For simplicity we consider here the case of a single subgroup and refer to Section 2.1 for the general case. We say that X is a *relative generating set* of G with respect to H, if G is generated by the set $H \cup X$.

In this situation there exists a canonical homomorphism

$$\varepsilon : F = H * F(X) \to G,$$

where $F(X)$ is the free group with the basis X. If $Ker\ \varepsilon$ is a normal closure of a subset $\mathcal{R} \subseteq N$ in F, we say that G has the *relative presentation*

$$\langle X, H \mid R = 1,\ R \in \mathcal{R} \rangle \tag{1.1}$$

The relative presentation (1.1) is *finite* if the sets X and $\mathcal{R}$ are finite. G is said to be *relatively finitely presented* with respect to H, if it admits a finite relative presentation. Note that G and H need not be finitely presented or even finitely generated in the usual sense.

Similarly one can define the notion of a relatively finitely generated and a relatively finitely presented group with respect to arbitrary collection of subgroups. We begin with the theorem, which shows some restrictions in case G is finitely generated (in the usual sense) and is finitely presented with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ (see Proposition 2.29 and Corollary 2.48).

Theorem 1.1. *Let G be a finitely generated group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. Suppose that G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$. Then the following conditions hold.*

1) The collection of subgroups is finite, i.e. card $\Lambda < \infty$.

2) Each subgroup H_λ is finitely generated.

Definition 1.2. Given a finite relative presentation (1.1), we say that $f : \mathbb{N} \to \mathbb{N}$ is *a relative isoperimetric function* of (1.1) if for every word W of length $\|W\| \le n$ over the alphabet $X^{\pm 1} \cup H$ representing the identity in G, there exists an expression

$$W =_F \prod_{i=1}^{k} f_i^{-1} R_i f_i \tag{1.2}$$

(with the equality in the group F), where $R_i \in \mathcal{R}$, $f_i \in F$, and $k \le f(n)$. The minimal relative isoperimetric function of (1.1) is called the *relative Dehn function* of G (with respect to H). We denote it by $\delta^{rel}_{G,H}$.

For instance, any finitely presented group G is relatively finitely presented with respect to the trivial subgroup. In this case the corresponding relative Dehn function coincides with the ordinary Denh function of G.

Example 1.3. We stress that the relative Dehn function is not always well–defined, i.e., it can be infinite for certain values of the argument, since the number of words of bounded relative length can be infinite. Indeed consider the group

$$G = \langle a, b \mid [a, b] = 1\rangle \cong \mathbb{Z} \times \mathbb{Z}$$

and the cyclic subgroup H generated by a. Clearly one can take $X = \{b\}$. It is easy to see that the word $W_n = [a^n, b]$ has length 4 as a word over $H \cup \{b\}$ for every n. However, the minimal number of factors in the expression of type (1.2) corresponding to W_n growths linearly as $n \to \infty$. Thus we have $\delta^{rel}_{G,H}(4) = \infty$ for this relative presentation.

The definition of the relative Dehn function in case of arbitrary collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ is similar (see Section 2.3). Analogously to the case of ordinary Dehn functions, if the relative Dehn function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ is finite for each value of the argument, it is independent of the choice of a finite relative presentation up to some equivalence relation (Theorem 2.34). Thus we can speak about the relative Dehn function of the pair $G, \{H_\lambda\}_{\lambda\in\Lambda}$ by means of the corresponding equivalence classes. The next result is obtained in Section 2.3 and provides us with necessary conditions for the relative Dehn function to be well–defined.

Theorem 1.4. *Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. Suppose that G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Denh function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$) is well–defined. Then the following conditions hold.*

1) For any $g_1, g_2 \in G$, the intersection $H_\lambda^{g_1} \cap H_\mu^{g_2}$ is finite whenever $\lambda \neq \mu$.

2) The intersection $H_\lambda^g \cap H_\lambda$ is finite for any $g \notin H_\lambda$.

The main reason for our study of relative Dehn functions is the following characterization of relative hyperbolicity of finitely generated groups.

Theorem 1.5. *Let G be a finitely generated group, $\{H_1, H_2, \ldots, H_m\}$ a collection of subgroups of G. Then the following conditions are equivalent.*

1) G is finitely presented with respect to $\{H_1, H_2, \ldots, H_m\}$ and the corresponding relative Dehn function is linear.

2) G is hyperbolic with respect to the collection $\{H_1, H_2, \ldots, H_m\}$ in the sense of Farb and satisfies the BCP property (or, equivalently, G is hyperbolic relative to $\{H_1, H_2, \ldots, H_m\}$ in the sense of Bowditch).

Theorem 1.5 allows to consider the definition below as a generalization of Bowditch's one.

Definition 1.6. We say that a group G is *hyperbolic relative to a collection of subgroups* $\{H_\lambda\}_{\lambda\in\Lambda}$ if G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Dehn function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ is linear.

Standard arguments show that if the relative Dehn function is subquadratic, then, in fact, it is linear (Corollary 2.57).

Given a group G, a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$, and a finite relative generating set X of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$, we denote by $\Gamma(G, X\cup\mathcal{H})$ the Cayley graph of G with respect to the generating set $X\cup\left(\bigcup\limits_{\lambda\in\Lambda} H_\lambda\right)$. It is easy to prove that if the relative Dehn function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ is linear, then $\Gamma(G, X\cup\mathcal{H})$ is hyperbolic. A partially converse result is obtained in Section 2.5.

Theorem 1.7. *Suppose that a group G is finitely presented with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Dehn function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ the is well–defined. Then the following conditions are equivalent.*

1) The group G is hyperbolic relative to $\{H_\lambda\}_{\lambda\in\Lambda}$.

2) The Cayley graph $\Gamma(G, X\cup\mathcal{H})$ is a hyperbolic metric space.

If the group G is generated by a finite set X in the ordinary non–relative sense, we can think of the Cayley graph of G with respect to X, $\Gamma(G, X)$, as a subgraph of the graph $\Gamma(G, X\cup\mathcal{H})$ defined above. Obviously these graphs have the same set of vertices. Assuming the lengths of any edge of $\Gamma(G, X)$ and $\Gamma(G, X\cup\mathcal{H})$ to have length 1, we get two combinatorial metrics $dist_X$ and $dist_{X\cup\mathcal{H}}$ on $\Gamma(G, X)$ and $\Gamma(G, X\cup\mathcal{H})$ respectively. The proof of the Theorem 1.5 given in the appendix is based on some results about the geometry of the embedding of $\Gamma(G, X)$ into $\Gamma(G, X\cup\mathcal{H})$, which are proved in Chapter 3. Although the proofs of these facts take a significant part of our paper, the theorems seem to be too technical to formulate them here.

We also introduce and study the notion of relative quasi–convexity for subgroups of G in case G is finitely generated.

Definition 1.8. Let G be a group generated by a finite set X, $\{H_1, \ldots, H_m\}$ a collection of subgroups of G. A subgroup R of G is called *relatively quasi–convex* with respect to $\{H_1, \ldots, H_m\}$ if there exists a constant $\sigma > 0$ such that the following condition holds. Let f, g be two elements of R, and p an arbitrary geodesic path from f to g in $\Gamma(G, X\cup\mathcal{H})$. Then for any vertex $v\in p$ there exists a vertex $w\in R$ such that

$$dist_X(u, w)\le\sigma.$$

In case G is hyperbolic relative to $\{H_1, \ldots, H_m\}$, this notion is independent of the choice of the finite generating set of G (Proposition 4.10). A subgroup R is called *strongly relatively quasi–convex* if, in addition, the intersections $R\cap H_i^g$ are finite for all $i = 1, \ldots, m$, $g\in G$.

The next three theorems are the relative analogues of well–known facts about quasi–convex subgroups of ordinary hyperbolic groups (see Section 4.2). In these theorems we suppose G to be a finitely generated group hyperbolic relative to a collection of subgroups $\{H_1, \ldots, H_m\}$.

Theorem 1.9. *Let R a subgroup of G. Then the following conditions are equivalent.*

1) R is strongly relatively quasi–convex.

2) R is generated by a finite set Y and the natural embedding $(R, dist_Y) \to (G, dist_{X\cup\mathcal{H}})$ is a quasi–isometry.

As a corollary, we obtain

Theorem 1.10. *If R is a strongly relatively quasi–convex subgroup of G, then R is a hyperbolic group.*

Finally we show that the set of relatively quasi–convex subgroups is closed under intersections.

Theorem 1.11. *Let P and R be two relatively quasi–convex subgroups of G. Then $P \cap R$ is relatively quasi–convex.*

Let us mention some applications of the technique developed in this paper to the study of algebraic and algorithmic properties of relatively hyperbolic groups. We say that an element $g \in G$ is *hyperbolic* if g is not conjugate to an element of one of the subgroups H_λ, $\lambda \in \Lambda$. In the next two theorems and their corollaries we assume that G is an arbitrary (not necessary finitely generated) group that is relatively hyperbolic with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$. The proofs can be found in Sections 4.1, 4.3.

Theorem 1.12. *There exist only finitely many conjugacy classes of hyperbolic elements of finite order in G. In particular, the set of orders of hyperbolic elements is finite.*

Corollary 1.13. *If G is residually finite and all subgroups H_λ are torsion free, then G is virtually torsion free, that is G contains a torsion free subgroup of finite index.*

It is worth to notice that the assumption of residually finiteness is essential in the last corollary. Indeed in Section 4.1 we construct an example of a (finitely generated) group G hyperbolic relative to a torsion–free subgroup H such that G is not virtually torsion–free.

Theorem 1.14. *For any hyperbolic element $g \in G$ of infinite order, there exist $\lambda > 0$, $c \geq 0$ such that*

$$dist_{X\cup\mathcal{H}}(1, g^n) > \lambda|n| - c$$

for any $n \in \mathbb{N}$.

It follows from the proof of Theorem 1.14 that every cyclic subgroup generated by a hyperbolic element of infinite order has finite index in its centralizer. In other terms this was first proved by Tukia in [81]. We also show that the constant λ in Theorem 1.14 can be chosen independently of g.

Corollary 1.15. *If g is a hyperbolic element of G of infinite order and g^k is conjugate to g^l for some $k, l \in \mathbb{Z}$, then $k = \pm l$*

We note that the existence of an action of G on a hyperbolic graph such that all edge stabilizers are trivial and every vertex stabilizer is conjugate to H_i for some i (that is a 'weak' form of hyperbolicity of G) is not sufficient for the fulfilment of Theorems 1.12 and 1.14. The counterexamples are provided in Sections 4.1, 4.3.

It is known that the word and the conjugacy problems are decidable in a (finitely generated) group G hyperbolic with respect to $\{H_1, \ldots, H_m\}$ provided these problems are decidable for each H_i [20], [37]. Also if the conjugacy problem is solvable in $H_1, \ldots, H_m$, then given $g \in G$ and $i \in \{1, \ldots, m\}$, it is possible to decide whether g is conjugate to an element of H_i [20]. Some of these results are generalized in Sections 5.1, 5.2.

The application of our approach to the study of other algorithmic problems leads to the following theorem.

Theorem 1.16. *Let G be a group hyperbolic with respect to a collection of recursively presented subgroups $\{H_1, \ldots, H_m\}$. Then each of the algorithmic problems listed below is solvable in G whenever the word problem is solvable in each of the subgroups $H_1, \ldots, H_m$.*

1) The conjugacy problem for hyperbolic elements, that is, given two hyperbolic elements $f, g \in G$, to decide whether f and g are conjugate.

2) The order problem for hyperbolic elements, that is to calculate the order of a given hyperbolic element $g \in G$.

3) The root problem for hyperbolic elements, that is, given an hyperbolic element $g \in G$, to decide whether there exists a nontrivial root of g in G.

4) The power conjugacy problem for hyperbolic elements, that is to decide, for any two hyperbolic elements $g, f \in G$, whether or not there exist two hyperbolic conjugate powers of f and g.

Acknowledgements. The paper was written in part during the author's visits in Courant Institute of Mathematical Sciences, in spring 2002, and Vanderbilt University, in fall 2002. I would like to thank Brian Bowditch, Benson Farb, Mihail Gromov, Alexander Ol'shanskii, and Mark Sapir for useful conversations and remarks.

Chapter 2

Relative isoperimetric inequalities

2.1 Relative presentations and length functions

We begin by introducing relative generating sets of groups with respect to fixed collection of subgroups.

Definition 2.1. Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G, X a subset of G. We say that G is *generated by* X *with respect to* $\{H_\lambda\}_{\lambda\in\Lambda}$ (or, equivalently, X is a *relative generating set of* G *with respect to* $\{H_\lambda\}_{\lambda\in\Lambda}$) if G is generated by the set $\left(\bigcup\limits_{\lambda\in\Lambda} H_\lambda\right)\cup X$. We will always assume the set X to be symmetrized, i.e., $X = X^{-1}$.

In the above situation the group G can be regarded as the quotient group of the free product

$$F = \left(\ast_{\lambda\in\Lambda}\widetilde{H}_\lambda\right) \ast F(X), \tag{2.1}$$

where the groups $\widetilde{H}_\lambda$ are isomorphic copies of H_λ, and $F(X)$ is the free group with the basis X. Let us denote by $\mathcal{H}$ the disjoint union

$$\mathcal{H} = \bigsqcup_{\lambda\in\Lambda}(\widetilde{H}_\lambda \setminus \{1\}).$$

It is easy to see that F is generated by $X\cup\mathcal{H}$.

Conventions and notation. By $(X\cup\mathcal{H})^*$, we denote the free monoid generated by $X\cup\mathcal{H}$. Given $W\in(X\cup\mathcal{H})^*$, $\|W\|$ denotes the length of the word W, $\overline{W}$ denotes the element of G represented by W. Throughout this paper we write $U\equiv V$ to express letter–for–letter equality of two words $U, V\in(X\cup\mathcal{H})^*$ and write $U =_F V$ when U and V represent the same elements of the group F.

To simplify our notation we identify words from $(X \cup \mathcal{H})^*$ and elements of the group F represented by them. We also write x^t for $t^{-1}xt$ and $[x,t]$ for $x^{-1}t^{-1}xt$.

For every $\lambda \in \Lambda$, we denote by $\mathcal{S}_\lambda$ the set of all words over the alphabet $\widetilde{H}_\lambda \setminus \{1\}$ that represent the identity in F. Thus the group F can be defined by the presentation

$$\langle X, \mathcal{H} \mid S = 1, S \in \bigcup_{\lambda \in \Lambda} \mathcal{S}_\lambda \rangle. \tag{2.2}$$

The isomorphisms $\widetilde{H}_\lambda$ and the identity map on X can be uniquely extended to a homomorphism $\epsilon : F \to G$. We denote its kernel by N.

Definition 2.2 (Relative presentation). We say that the group G has the *relative presentation*

$$\langle X, \mathcal{H} \mid S = 1, S \in \bigcup_{\lambda \in \Lambda} \mathcal{S}_\lambda, R = 1, R \in \mathcal{R} \rangle, \tag{2.3}$$

with respect to the collection of subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$, where $\mathcal{R} \subseteq (X \cup \mathcal{H})^*$, if N is the normal closure of the set $\mathcal{R}$ in the group F. It is convenient to assume that $\mathcal{R}$ is symmetrized that is for every $R \in \mathcal{R}$, the set $\mathcal{R}$ contains all cyclic shifts of R and R^{-1}.

For brevity, we use the following reduced record for the presentation (2.3)

$$\langle X, H_\lambda, \lambda \in \Lambda \mid R = 1, R \in \mathcal{R} \rangle, \tag{2.4}$$

Definition 2.3. The relative presentation (2.4) is called *finite* if both the sets $\mathcal{R}$ and X are finite. If there exists a finite relative presentation of a group G with respect to a collection of subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$, we say that G is *finitely presented* relative to $\{H_\lambda\}_{\lambda \in \Lambda}$.

Example 2.4. Consider the amalgamated product

$$G = H_1 *_{K=L} H_2$$

of two arbitrary groups H_1, H_2 associated to an isomorphism $\alpha : K \to L$ between subgroups $K \leq H_1$ and $L \leq H_2$. Then G has the relative presentation

$$\langle H_1, H_2 \mid k = \alpha(k),\ k \in K \rangle$$

with respect to $\{H_1, H_2\}$. If K is finitely generated, one can construct a finite relative presentation for G with respect to $\{H_1, H_2\}$. Indeed it is sufficient to impose the relations of type $k = \alpha(k)$ for all generators of K.

Example 2.5. If H is a group and $\alpha : A \to B$ is an isomorphism between two subgroups $A, B \leq H$, then the corresponding HNN–extension has the relative presentation

$$\langle H \mid a^t = \alpha(a), a \in A \rangle.$$

As above this relative presentation can be made finite in case A is finitely generated.

Definition 2.6. To each element $g \in G$, we assign its *relative length with respect to the collection of subgroups* $\{H_\lambda\}_{\lambda\in\Lambda}$ of G (or simply *relative length*) $|g|_{X\cup\mathcal{H}}$ that is the length of a shortest word from $(X \cup \mathcal{H})^*$ representing g in G. This length function induces the left–invariant *relative distance function* on $G \times G$ by the rule

$$dist_{X\cup\mathcal{H}}(g, h) = |g^{-1}h|_{X\cup\mathcal{H}}.$$

We note that the group G endowed with the relative metric is not always a proper metric space, i.e., the closed balls can be infinite. It is easy to see that G is proper if and only if $card\,\Lambda < \infty$ and all subgroups H_λ are finite.

It is clear that this distance strongly depends on the choice of the set X and the collection of subgroups. However, if the collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ is fixed and G is finitely generated with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$, then the asymptotic behavior of the distance function is essentially independent of the choice of a finite relative generating set. This can be expressed in the following way.

Definition 2.7. Two metrics $dist_1$ and $dist_2$ on the same space are called *Lipschitz equivalent* if the ratios $dist_1/dist_2$ and $dist_2/dist_1$ are bounded when they are considered as functions on the Cartesian square of the space minus the diagonal.

The proposition below is the relative analogue of the well known property of ordinary word metrics on finitely generated groups.

Proposition 2.8. *Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. Suppose that X and Y are two finite relative generating sets of G with respect to $\{H_\lambda\}_\Lambda$. Then the corresponding distance functions $dist_{X\cup\mathcal{H}}$ and $dist_{Y\cup\mathcal{H}}$ are Lipschitz equivalent.*

Proof. Let us take an arbitrary element $g \in G$. Suppose that $W \in (Y \cup \mathcal{H})^*$ is a shortest word representing g. For every $y \in Y$, we fix a word $V_y \in (X \cup \mathcal{H})^*$ that represents y. Put

$$M = \max_{y\in Y} \|V_y\|.$$

As the set Y is finite, we have $M < \infty$.

If we consider the word $U \in (X \cup \mathcal{H})^*$ that is obtained from W by replacing all letters $y \in Y$ with the corresponding words V_y, then

$$\|U\| \le M\|W\| = M|g|_{Y\cup\mathcal{H}}.$$

Finally we have

$$|g|_{X\cup\mathcal{H}} \le \|U\| \le M|g|_{Y\cup\mathcal{H}}.$$

The reverse inequality can be obtained in the analogous way. The proposition is proved. □

2.2 Geometry of van Kampen diagrams over relative presentations

Our study of properties of relatively hyperbolic groups is based on the combinatorial geometry of van Kampen diagrams over relative representations. In this section we collect definitions and some technical facts about the diagrams which are used in what follows.

Definition 2.9. Recall that a planar *map* Δ over a presentation

$$\langle Z \mid \mathcal{P} \rangle \tag{2.5}$$

is a finite oriented connected simply–connected 2–complex endowed with a labelling function $\phi : E(\Delta) \to Z^{\pm 1}$, where $E(\Delta)$ denotes the set of oriented edges of Δ, such that $\phi(e^{-1}) = (\phi(e))^{-1}$. The label of a path $p = e_1 \ldots e_n$ is, by definition, the word $\phi(e_1) \ldots \phi(e_n)$. By *length* $l(p)$ we mean the number of edges in p. Given a cell Π of Δ, we denote by $\partial\Pi$ the boundary of Π; similarly, $\partial\Delta$ denotes the boundary of Δ. The labels of $\partial\Pi$ and $\partial\Delta$ are defined up to a cyclic permutation.

Definition 2.10. A map Δ over a presentation (2.5) is called a *van Kampen diagram* over (2.5) if the following holds. For any cell Π of Δ, the boundary label $\phi(\partial\Pi)$ is equal to a cyclic permutation of a word $P^{\pm 1}$, where $P \in \mathcal{P}$. Sometimes it is convenient to use the notion of 0–refinement in order to assume diagrams to be homeomorphic to a disc. We do not explain here this notion and refer the interested reader to [61, Ch. 4].

The van Kampen lemma states that a word W over the alphabet $Z^{\pm 1}$ represents the identity in the group given by (2.5) if and only if there exists a simply–connected planar diagram Δ over (2.5) such that $\phi(\partial\Delta) \equiv W$ [56].

Dealing with diagrams over relative presentations we will divide the set of 2–cells into two parts as follows.

Definition 2.11. Let Δ be a van Kampen diagram over the relative presentation (2.3). A cell Π of Δ is called an $\mathcal{S}_\lambda$*–cell* if it corresponds to a relator from $\mathcal{S}_\lambda$. A cell Π is called an $\mathcal{S}$–cell if it is an $\mathcal{S}_\lambda$–cell for some $\lambda \in \Lambda$. Obviously for any $\mathcal{S}$–cell, the label of its boundary represents 1 in the group F. Similarly we call Π an $\mathcal{R}$–cell if it corresponds to a relator from $\mathcal{R}$.

Definition 2.12. Given a van Kampen diagram Δ over (2.3), we denote by $N_{\mathcal{R}}(\Delta)$ (respectively by $N_{\mathcal{S}}(\Delta)$) the number of $\mathcal{R}$–cells (respectively $\mathcal{S}$–cells) of Δ. We define the area of Δ by the formula

$$Area(\Delta) = N_{\mathcal{R}}(\Delta) + N_{\mathcal{S}}(\Delta).$$

Definition 2.13. The *type* of the diagram Δ is the triple $\tau(\Delta) = (N_{\mathcal{R}}(\Delta), N_{\mathcal{S}}(\Delta), card\, E(\Delta))$. We fix the lexicographic order on the set of all



Figure 2.1: Types of edges in diagrams.

triples, that is,

$$(a_1, b_1, c_1) \le (a_2, b_2, c_2) \Longleftrightarrow$$

$$(a_1 < a_2) \ \vee\ (a_1 = a_2 \ \wedge\ b_1 < b_2) \ \vee\ (a_1 = a_2 \ \wedge\ b_1 = b_2 \ \wedge\ c_1 \le c_2).$$

We will exploit the following classification of edges of van Kampen diagrams over group presentations.

Definition 2.14. A (non–oriented) edge e of a diagram Δ is called *internal* if it is a common edge of two cells of Δ; otherwise e is called *external.* Thus the union of all external edges forms the boundary $\partial\Delta$ of the diagram Δ. Further an external edge e is called an *external edge of the first type* if e belongs to the boundary of some cell of the diagram; otherwise e is called an *external edge* of the second type.

The notions introduced in Definition 2.14 are illustrated on Fig. 2.1. The edges $e_1, \ldots, e_5$ are external, the edge d is internal. More precisely, $e_1, \ldots, e_4$ are external edges of the first type and e_5 is external of the second type.

Lemma 2.15. *Let Δ be a van Kampen diagram over (2.3). Suppose that Δ has the minimal type among all van Kampen diagrams over (2.3) with the same boundary label. Then every internal edge of Δ belongs to the boundary of some $\mathcal{R}$–cell of Δ.*

Proof. Let $e \in E(\Delta)$ be an internal edge. Suppose that e does not belong to the boundary of any $\mathcal{R}$–cell of Δ. Then there are two possibilities (see Fig. 2.2): either e is a common edge of two distinct S_λ–cells Π_1 and Π_2 or there is an S_λ–cell Π in Δ such that

$$\partial\Pi = ed_1e^{-1}d_2,$$

where d_1 and d_2 are cycles in Δ (one of them may be trivial) and one of the cycles d_1, d_2, say d_2, is contained in the part of the diagram bounded by the other one. In the first case let

$$\partial\Pi_1 = ec_1, \qquad \partial\Pi_2 = c_2^{-1}e^{-1}.$$

Note that the label of the path $c_1c_2^{-1}$ (respectively d_1) consists of letters from $\widetilde{H}_\lambda$ and represents 1 in F. Since $\mathcal{S}_\lambda$ contains all words over $\widetilde{H}_\lambda$ that represents



Figure 2.2: Two cases in the proof of Lemma 2.15.

the identity in the group F, we can replace the cells Π_1, Π_2 (respectively the cell Π) with the cell Σ with the boundary $c_1c_2^{-1}$ (respectively d_1). In both cases the type of the diagram decreases and we get a contradiction. $\square$

Corollary 2.16. *Let Δ be a van Kampen diagram over (2.3). By $S(\Delta)$ we denote the set of all $\mathcal{S}$–cells of Δ. Suppose that (2.3) is relatively finite and Δ has the minimal type among all van Kampen diagrams over (2.3) with the same boundary label. Then we have*

$$\sum_{\Pi\in S(\Delta)} l(\partial\Pi) \le MN_{\mathcal{R}}(\Delta) + l(\partial\Delta), \tag{2.6}$$

where $M = \max\limits_{R\in\mathcal{R}} \|R\|$.

Proof. Let $R(\Delta)$ denote the set of all $\mathcal{R}$–cells of Δ. According to Lemma 2.15, the number of all (non–oriented) edges of Δ satisfies the inequality

$$card\ E(\Delta) \le \sum_{\Xi\in R(\Delta)} l(\partial\Xi) + l(\partial\Delta) \le MN_{\mathcal{R}}(\Delta) + l(\partial\Delta).$$

This implies (2.6) as every edge of Δ belongs to the boundary of at most one $\mathcal{S}$–cell. $\square$

Definition 2.17. The *Cayley graph* $\Gamma = \Gamma(G, Z)$ of a group G generated by a set Z is an oriented labelled 1–complex with the vertex set $V(\Gamma) = G$ and the edge set $E(\Gamma) = G \times Z$. An edge $e = (g, s) \in E(\Gamma)$ goes from the vertex g to the vertex gs and has the label $\phi(e) = s$. As usual, we denote the origin and the terminus of the edge e, i.e., the vertices g and gs, by e_- and e_+ respectively.

Given a combinatorial path $p = e_1e_2\ldots e_k$ in Γ, where $e_1, e_2, \ldots, e_k \in E(\Gamma)$, we denote by $\phi(p)$ its label. As in the case of diagrams, $\phi(p) = \phi(e_1)\phi(e_2)\ldots\phi(e_k)$. By $p_- = (e_1)_-$ and $p_+ = (e_k)_+$ we denote the origin and the terminus of p respectively. A path p is called *irreducible* if it contains no subpaths of type ee^{-1} for $e \in E(\Gamma)$. A *subpath* q of $p = e_1e_2\ldots e_k$ is a path of type $e_ie_{i+1}\ldots e_j$ for some $1 \le i \le j \le k$. (So we always assume p and q to have compatible orientations, that is, starting from p_- and passing along p, we first meet the vertex q_- and then q_+.) The graph Γ can be regarded as a metric space if we endow it with a combinatorial metric. This means that the length of every edge of Γ is assumed to be equal to 1.

In this paper we denote by $\Gamma(G, X \cup \mathcal{H})$ the Cayley graph of the group G given by (2.3) with respect to the generating set $X \cup \mathcal{H}$. Note that every van Kampen diagram Δ over (2.3) can be mapped to $\Gamma(G, X \cup \mathcal{H})$ in such a way that the mapping preserves labels and orientation. Taking into account this remark, we will often consider configurations in the Cayley graph $\Gamma(G, X \cup \mathcal{H})$ instead of the treating the corresponding van Kampen diagrams.

Now we are going to introduce an auxiliary terminology, which plays an important role in our paper.

Definition 2.18 (H_λ–subwords). Given a word $W \in (X \cup \mathcal{H})^*$, we say that a subword V of W is an H_λ*–subword* if V consists of letters from $\widetilde{H}_\lambda$. An H_λ–subword of W is called an H_λ*–syllable* if it is maximal, i.e., it is not contained in a bigger H_λ–subword of W.

As usual, by a cyclic word W we mean the set of all cyclic shifts of W. As in the case of ordinary words, we say that subword V of a cyclic word W is a H_λ–subword if it is an H_λ–subword of a certain cyclic shift of W. A maximal H_λ–subword of a cyclic word W is called an H_λ–syllable.

Definition 2.19 (H_λ–components). Let q be a path (respectively cyclic path) in $\Gamma(G, X \cup \mathcal{H})$. A subpath p of q is called an H_λ*–subpath*, if the label of p is an H_λ–subword of the word $\phi(q)$ (respectively cyclic word $\phi(q)$). A *component* (or more precisely an H_λ*–component*) of q is an H_λ–subpath p such that the label of p is an H_λ–syllable of the the word $\phi(q)$ (respectively cyclic word $\phi(q)$).

Definition 2.20 (Connected components). Two H_λ–components p_1, p_2 of a path q (cyclic or not) in $\Gamma(G, X \cup \mathcal{H})$ are called *connected* if there exists a path c in $\Gamma(G, X \cup \mathcal{H})$ that connects some vertex of p_1 to some vertex of p_2 and $\phi(c)$ is a word consisting of letters from $\widetilde{H}_\lambda$. The path c is called an H_λ*–connector.* Note that this is equivalent to the requirement that for any two vertices v_1 and v_2 of p_1 and p_2 respectively there exists a connector c such that $c_- = v_1, c_+ = v_2$. (In algebraic terms this means that these two vertices belong to the same coset gH_λ.) Clearly we can always assume that c consists of a single edge, as every element of H_λ is included in the set of generators.

Sometimes we will also speak about connected H_λ–syllables in a word $W \in (X \cup \mathcal{H})^*$ (cyclic or not). By these we mean two H_λ–syllables U, V of W such that the corresponding components of some (or, equivalently, of any) path in $\Gamma(G, X \cup \mathcal{H})$ labelled W are connected.

Definition 2.21 (Isolated components). An H_λ–component p of a path q (cyclic or not) is called *isolated* if no (distinct) H_λ–component is connected to p. The notion of an isolated H_λ–syllable of a word $W \in (X \cup \mathcal{H})^*$ is defined in the obvious way.



Figure 2.3: The path corresponding to the word $W \equiv a^2 a^t a^3$.

Example 2.22. Let us consider the Baumslag–Solitar group

$$BS(1,2) = \langle a, t \mid a^t = a^2 \rangle.$$

Set $H = \langle a \rangle$ and consider the word $W \equiv a^2 a^t a^3$. Then the H–syllables a^3 and a^2 are connected since a^t represents the same element of G as a^2 (see Fig. 2.3). The H-syllable a of W is isolated.

In the next sections we will often use the following result without any references. The proof follows from the definitions in the obvious way.

Lemma 2.23. *Let p be a geodesic path in the Cayley graph $\Gamma(G, X \cup \mathcal{H})$. Then every component of p is isolated.*

Definition 2.24. A relative representation (2.3) of a group G with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ is called *reduced* if each relator $R \in \mathcal{R}$ has minimal length among all words from $(X \cup \mathcal{H})^*$ representing the same element of the group F. In particular this means that for any $\lambda \in \Lambda$ and any $R \in \mathcal{R}$, every H_λ–syllable of R consists of a single letter. Without loss of generality we may assume all relative presentations under consideration to be reduced.

The next lemma shows that, without loss of generality, we can assume finite relative presentations to be reduced.

Definition 2.25. For every $\lambda \in \Lambda$, we denote by Ω_λ the subset of all elements $g \in H_\lambda$ such that there exists a relator $R \in \mathcal{R}$, and an H_λ–syllable V of R such that V represents g in G. We also put

$$\Omega = \bigcup_{\lambda\in\Lambda} \Omega_\lambda.$$

It is important that the set Ω is finite, whenever the set $\mathcal{R}$ is finite.

Definition 2.26 (Relative area of a cycle in the Cayley graph). Let q be a cycle in the Cayley graph $\Gamma(G, X \cup \mathcal{H})$. We define its *relative area*, $Area^{rel}(q)$, with respect to the presentation (2.3) as the minimal number $N_{\mathcal{R}}(\Delta)$ of $\mathcal{R}$–cells among all van Kampen diagrams Δ over (2.3) with the boundary label $\phi(\partial\Delta) \equiv \phi(q)$.

Now we are ready to formulate the main lemma of this section. Although Lemma 2.27 is quite trivial, it allows to obtain some important results on groups



Figure 2.4: Two cases in the proof of Lemma 2.27; the component p_i' is marked by the thick line.

given by finite relative presentations (see, for example, Proposition 2.29 and Theorem 2.36 below). Recall that for a word $W \in (X \cup \mathcal{H})^*$, $\overline{W}$ means the element of G represented by W.

Lemma 2.27. *Suppose that a group G is given by the reduced finite relative presentation (2.3) with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$. Let q be a cycle in $\Gamma(G, X\cup\mathcal{H})$, $p_1, \ldots, p_k$ a certain set of isolated H_λ–components of q. Then*

$$\overline{\phi(p_i)} \in \langle \Omega_\lambda \rangle \tag{2.7}$$

for any $i = 1, \ldots, k$. Moreover, the lengths of the elements $\overline{\phi(p_1)}, \ldots, \overline{\phi(p_k)}$ with respect to the generating set Ω_λ of the subgroup $\langle\Omega_\lambda\rangle$ satisfy the inequality

$$\sum_{i=1}^{k} |\overline{\phi(p_i)}|_{\Omega_\lambda} \le M \cdot Area^{rel}(q), \tag{2.8}$$

where

$$M = \max_{R\in\mathcal{R}} ||R||.$$

Proof. Changing the enumeration of the components $p_1, p_2, \ldots, p_k$ if necessarily, we may assume that $q = p_1q_1p_2q_2\ldots p_kq_k$ for some paths $q_1, \ldots, q_k$ in $\Gamma(G, X\cup$

$\mathcal{H}$). By $\mathcal{Q}$ we denote the set of all irreducible cycles q' with the same initial point as q in $\Gamma(G, X \cup \mathcal{H})$ such that q' can be represented as

$$q' = p'_1 q_1 p'_2 q_2 \ldots p'_k q_k, \tag{2.9}$$

where $p'_1, p'_2, \ldots, p'_n$ are some H_λ–components of q' satisfying the conditions $\overline{(p'_i)_-} = \overline{(p_i)_-}$ and $(p'_i)_+ = (p_i)_+$ for $i = 1, 2, \ldots, k$. In particular, we have $\overline{\phi(p'_i)} = \overline{\phi(p_i)}$ for any $i = 1, \ldots, k$. Note that for any $q' \in \mathcal{Q}$, the subpaths $p'_1, p'_2, \ldots, p'_k$ are isolated H_λ–components of q'. By $\mathcal{D}$ we denote the set of all diagrams Δ over (2.3) such that $\phi(\partial\Delta) \equiv \phi(q')$ for some $q' \in \mathcal{Q}$.

Now let Δ be a diagram of minimal type in $\mathcal{D}$. To simplify our notation we will identify the boundary of Δ with the corresponding cycle $q' = p'_1 q_1 p'_2 q_2 \ldots p'_k q_k$ in $\Gamma(G, X \cup \mathcal{H})$. We are going to show that every edge of subpaths $p'_1, p'_2, \ldots, p'_k$ of $\partial\Delta$ belongs to the boundary of some $\mathcal{R}$–cell.

Suppose this is not true; then there are two possibilities: either a certain p'_i contains an external edge of the second type, or at least one $\mathcal{S}$–cell has a common edge with p'_i.

Case 1. First assume that a certain component p'_i, contains an external edge d of the second type. Then $\partial\Delta = dc_2d^{-1}c_1$. Since p'_i is isolated, the cycle dc_2d^{-1} is a subpath of p'_i (see Fig. 2.4). As q' is irreducible, the subdiagram Σ bounded by the cycle dc_2d^{-1} contains at least one cell. However, this contradicts to our choice of Δ as we can decrease the type of the diagram by eliminating the subdiagram Σ.

Case 2. Now assume that some $\mathcal{S}$–cell Π has a common edge with p'_i for some i. Obviously Π is an S_λ–cell. Let $\partial\Pi = e^{-1}r$, where e is the largest common subpath of p'_i and $\partial\Pi$, and let $\partial\Delta = tp'_i$, where $p'_i = s_2es_1$ (see Fig. 2.4). Then we can decrease the type of the diagram by passing to the subdiagram bounded by the paths ts_2rs_1, which obviously belongs to $\mathcal{Q}$. Thus we get a contradiction again.

Therefore, each edge of any subpath p'_i of $\partial\Delta$ belongs to the boundary of some $\mathcal{R}$–cell of Δ. Since we assumed (2.3) to be reduced, the length of any H_λ–component of any $R \in \mathcal{R}$ equals 1 and so the element represented by the label of any edge of any p'_i belongs to Ω_λ. This gives (2.7). Finally, by our choice of Δ, we have $N_\mathcal{R}(\Delta) \le Area^{rel}(q)$. Hence the total number of edges of $p'_1, p'_2, \ldots, p'_k$ satisfies the inequality

$$\sum_{i=1}^{k} l(p'_i) \le MN_\mathcal{R}(\Delta) \le MArea^{rel}(q),$$

which yields (2.8). The lemma is proved. □

Note that for any path q in $\Gamma(G, X \cup \mathcal{H})$, every H_λ–component p of q is contained in the unique maximal system of connected H_λ–components of q. In the next section we will use the following corollary of Lemma 2.27.



Figure 2.5: The decomposition of the cycle q in the proof of Corollary 2.28.

Corollary 2.28. *Suppose that a group G is given by the reduced finite relative presentation (2.3). Let $q = pr$ be a cycle in the Cayley graph $\Gamma(G, X \cup \mathcal{H})$, where p an H_λ–component of q for some $\lambda \in \Lambda$. Let $p, p_1, p_2, \ldots, p_l$ be the corresponding maximal system of connected H_λ–components of q. Then*

$$\overline{\phi(p)} \in \langle \overline{\phi(p_1)}, \overline{\phi(p_2)}, \ldots, \overline{\phi(p_l)}, \Omega_\lambda \rangle. \tag{2.10}$$

Proof. Without loss of generality we may assume that

$$r = r_1 p_1 \ldots r_l p_l r_{l+1}$$

for some paths $r_1, \ldots, r_l + 1$. Let $c_1, \ldots, c_{l+1}$ be edges in $\Gamma(G, X \cup \mathcal{H})$ labelled by elements of $\widetilde{H}_\lambda$ such that

$$(c_1)_- = p_+, \quad (c_{l+1})_+ = p_-,$$

and

$$(c_i)_+ = (p_i)_-, \quad (c_{i+1})_- = (p_i)_+$$

for $i = 1, \ldots, l$ (see Fig. 2.5).

Let us consider cycles $q_j = c_j r_j^{-1}$. Since the system $p, p_1, \ldots, p_l$ is maximal, it follows that c_j is an isolated H_i–component of q_j for every j. Applying Lemma 2.27 for cycles q_j, $j = 1, \ldots, l + 1$, we obtain

$$\overline{\phi(c_j)} \in \langle \Omega_\lambda \rangle.$$

This immediately implies (2.10) since $p = (c_1 p_1 \ldots c_l p_l c_{l+1})^{-1}$. ☐

Applying Lemma 2.27, we can prove the following.

Proposition 2.29. *Let G be a group, $\{H_\lambda\}_{\lambda \in \Lambda}$ a collection of subgroups of G. Suppose that G is generated by a finite set X in the ordinary non–relative sense and is finitely presented with respect to $\{H_\lambda\}_{\lambda \in \Lambda}$. Then each subgroup H_λ is generated by Ω_λ. In particular, H_λ is finitely generated.*

Proof. Suppose that G is given by a reduced finite relative presentation (2.3). Let us fix $\lambda \in \Lambda$. For every nontrivial element $h \in H_\lambda$, there is a word W over X such that $\overline{W} = h$ in G. Let us consider a cycle $q = pr$ in $\Gamma(G, X \cup \mathcal{H})$ such that $\overline{\phi(p)} = h$ and $\phi(r) \equiv W^{-1}$. Clearly p is an isolated H_λ–component of q as r contains no edges labelled by elements of $\widetilde{H}_\lambda$. Applying Lemma 2.27 for $k = 1$, $p_1 = p$, we obtain $h \in \langle \Omega_\lambda \rangle$. Therefore Ω_λ generates H_λ. As the presentation is finite, the set Ω_λ is finite and the proposition is proved. □

2.3 Relative Dehn functions

In the case of a finitely generated group G and a finitely generated subgroup $H \leq G$, the relative Dehn function corresponding to the pair (G, H) was introduced in [17] as a special case of the Howie function of a complex of groups (see [16] and [29] for details). However, the straightforward extension of the definition from [17] does not work in case the number of subgroups is greater that one.

Indeed, the definition from [17] involves the following construction. Let (L, K) be a geometric realization of the pair (G, H), i.e., a pair of connected combinatorial 2–complexes with finite 1–skeletons such that K is a subcomplex of L and there are finitely many cells in $L \setminus K$. Moreover, suppose that for any choice of base point in K, there exist isomorphisms $\pi_1(L) \to G$, $\pi_1(K) \to H$ for which the following diagram is commutative

$$\begin{tikzcd}
\pi_1(L) \arrow[r] & G \\
\pi_1(K) \arrow[r] \arrow[u] & H \arrow[u]
\end{tikzcd}$$

(The vertical maps are the natural embeddings.) Then one can form the combinatorial quotient $Q = L/K$ by contracting K to a point (see [41]). The relative Dehn function of the pair (G, H) is defined as the Dehn function of the universal cover E of Q related to the associated complex of groups ([17], [29]).

However, if $L = K_1 \cup K_2$, we obtain a point by contracting both K_1 and K_2. This situation arises when we consider a free product $G = H_1 * H_2$ of two finitely presented groups H_1 and H_2, and the corresponding two–complexes L, K_1, K_2 canonically associated with the finite presentations of G, H_1, and H_2 respectively.

In the present paper we exploit another approach based on the notion of relative presentation. Our treatment is inspired by the paper [70] and is similar to that in [17]. However we use a language, which is rather combinatorial in contract to geometric one in [17]. This allows to define the notion of the relative Dehn function with respect to an arbitrary set of arbitrary, not necessarily finitely generated, subgroups. In the case of a single subgroup, our definition is equivalent to that of Brick and Corson.

Suppose that G is given by the relative presentation 2.3. For a word $W \in$

$(X \cup \mathcal{H})^*$ such that W represents 1 in G, there exists an expression

$$W =_F \prod_{i=1}^{k} f_i^{-1} R_i f_i \tag{2.11}$$

with the equality in the group F given by 2.1, where $R_i \in \mathcal{R}$ and $f_i \in F$ for any i.

Definition 2.30. We say that a function $f : \mathbb{N} \to \mathbb{N}$ is a *relative isoperimetric function* of the presentation (2.3) with respect to subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$ if for any $n \in \mathbb{N}$ and any word $W \in (X \cup \mathcal{H})^*$ of length $\|W\| \leq n$ representing the identity in the group G, one can write W as in (2.11) with $k \leq f(n)$.

Notice that the above definition coincides with the definition of the ordinary Dehn function in case $\mathcal{H} = \emptyset$. The standard arguments show that Definition 2.30 is equivalent to the following.

Definition 2.31. A function $f : \mathbb{N} \to \mathbb{N}$ is a *relative isoperimetric function* of the presentation (2.3) with respect to subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$ if for any $n \in \mathbb{N}$ and any cycle q in the Cayley graph $\Gamma(G, X \cup \mathcal{H})$ of length $l(q) \leq n$, the relative area of q with respect to the presentation (2.3) satisfies

$$Area^{rel}(q) \leq f(n).$$

Definition 2.32. The smallest relative isoperimetric function of (2.3) is called the *relative Dehn function* of (2.3) with respect to subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$. As was notified in the introduction, it can happen that (2.3) does not posses any finite relative isoperimetric function. In this case we say that the relative Dehn function is not well–defined.

As in the usual (non–relative) case, we consider the relative isoperimetric functions up to the following equivalence relation.

Definition 2.33. For two functions $f, g : \mathbb{N} \to \mathbb{N}$, we say that f is asymptotically less than g and write $f \preceq g$ if there exist constants $C, K, L \in \mathbb{N}$ such that

$$f(n) \leq Cg(Kn) + Ln.$$

Further, we say that f is asymptotically equivalent to g and write $f \sim g$ if $f \preceq g$ and $g \preceq f$.

Theorem 2.34. *Let*

$$\langle X_1, H_\lambda, \lambda \in \Lambda \mid R = 1, R \in \mathcal{R}_1 \rangle \tag{2.12}$$

and

$$\langle X_2, H_\lambda, \lambda \in \Lambda \mid R = 1, R \in \mathcal{R}_2 \rangle \tag{2.13}$$

be two finite relative presentations of the same group G with respect to a fixed collection of subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$, δ_1 and δ_2 the corresponding relative Dehn functions. Suppose that δ_1 is well–defined, i.e., $\delta_1(n)$ is finite for every n. Then δ_2 is well–defined and $\delta_1 \sim \delta_2$.

Proof. Although the presentations are infinite (in the usual, non–relative sense), passing from (2.12) to (2.13) we change only a finite part of the presentation as X_i and $\mathcal{R}_i$ are finite for $i = 1, 2$. Thus in order to prove our theorem we have to repeat word–for–word the proof of its non–relative analogue replacing the word "length" with "relative length" everywhere. Proposition 2.8 will play the same role as the fact that the word metric on a finitely generated group is independent of the choice of a finite generating set up to the Lipschitz equivalence. This is straightforward and we leave details to the reader. $\square$

In what follows, speaking about the relative Dehn function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ we always mean the corresponding equivalence class.

Definition 2.35. Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. We say that G is *hyperbolic relative to* $\{H_\lambda\}_{\lambda\in\Lambda}$, if G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Denh function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda})$ is linear.

Proposition 2.36. *Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. Suppose that G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Denh function δ^{rel} of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda})$ is well–defined. Then the following conditions hold.*

1) The intersection $H_\lambda^{g_1}\cap H_\mu^{g_2}$ is finite whenever $\lambda\neq\mu$.

2) The intersection $H_\lambda^{g}\cap H_\lambda$ is finite for any $g\neq H_\lambda$.

Proof. First suppose that $\lambda\neq\mu$. It is sufficient to check that $H_\lambda^{g}\cap H_\mu$ is finite for every $g\in G$ since $H_\lambda^{g_1}\cap H_\mu^{g_2}=(H_\lambda^{g}\cap H_\mu)^{g_2}$ for $g=g_1g_2^{-1}$.

Consider a word $W\in(X\cup\mathcal{H})^*$ that represents g and has length

$$\|W\|=|g|_{X\cup\mathcal{H}}.$$

Assume that $W=W_1W_2$, where W_1 is the maximal (may be empty) prefix of W consisting of letters from $\widetilde{H}_\lambda$. Denote by f the element of G represented by W_2. It is clear that $H_\lambda^{g}=H_\lambda^{f}$. Thus it suffices to show that $H_\lambda^{f}\cap H_\mu$ is finite.

Taking into account this remark, we can always assume that if $g\neq 1$, then the first letter of the shortest word $W\in(X\cup\mathcal{H})^*$ representing the element g does not belong to H_λ.

Let us take an arbitrary element $h\in H_\lambda^{g}\cap H_\mu$ and denote by h_1, h_2 the letters from $\widetilde{H}_\lambda$ and $\widetilde{H}_\mu$ that represent elements $h^{g^{-1}}\in H_\lambda$ and $h\in H_\mu$ respectively. Since

$$\overline{W^{-1}h_1W}=\overline{h_2},$$

there is a cycle q in $\Gamma(G, X\cup\mathcal{H})$ having the label

$$\phi(q)\equiv W^{-1}h_1Wh_2^{-1},$$

of relative area

$$Area^{rel}(q)\leq\delta^{rel}(l(q))\leq\delta^{rel}(2\|W\|+2)=\delta^{rel}(2|g|_{X\cup\mathcal{H}}+2). \tag{2.14}$$



Figure 2.6: The cycle in $\Gamma(G, X \cup \mathcal{H})$ associated with $h \in H_\lambda^g \cap H_\mu$.

Let

$$q = v_1 p_1 v_2 p_2, \tag{2.15}$$

where

$$\phi(v_1) \equiv W^{-1}, \quad \phi(v_2) \equiv W, \tag{2.16}$$

and

$$\phi(p_1) = h_1, \quad \phi(p_2) = h_2^{-1}. \tag{2.17}$$

Note that the subpath p_1 is an isolated H_λ–component of q. Indeed, since the first letter of W does not belong to H_λ, p_1 is an H_λ–component. Suppose that there is another H_λ–component p of q that is connected to p_1. Since $\lambda \neq \mu$, p is a subpath of v_1 or v_2. For definiteness, assume that

$$v_1 = v_1' p v_2'',$$

where $|v_1''| \neq 0$, and there exists a path c in $\Gamma(G, X \cup \mathcal{H})$ that connects some vertex of p to a vertex of p_1 and has label consisting of letters from $\widetilde{H}_\lambda$ (see Fig. 2.6). Then $\phi(v_1'')$ represents an element of H_λ in G. Hence we can take a letter $k \in \widetilde{H}_\lambda$ such that

$$\overline{k} = \overline{\phi(p)\phi(v_2'')}$$

and consider the word $\phi(v_1')k \in (X \cup \mathcal{H})^*$, which is shorter than W and represents the same element g of G. A contradiction.

Thus p_1 is an isolated H_λ–component of q and we can apply Lemma 2.27. Using (2.14), we obtain

$$|h_1|_{\Omega_\lambda} = |\overline{\phi(p_1)}|_{\Omega_\lambda} \leq M \cdot Area^{rel}(q) \leq M\delta^{rel}(2|g|_{X \cup \mathcal{H}} + 2). \tag{2.18}$$

If $\delta^{rel}(n)$ is finite for any $n \in \mathbb{N}$, this means that the length of every element $h_1 \in H_\lambda^g \cup H_\mu$ with respect to Ω_λ is bounded by a constant which is independent of h_1. As Ω_λ is finite, we have $|H_\lambda^g \cap H_\mu| < \infty$.

Let us prove the second assertion of the theorem. Suppose that $g \notin H_\lambda$ and W is as above. Arguing in the analogous way, we can assume that the first letter of the word $W \in (X \cup \mathcal{H})^*$ does not belong to H_λ. As in the previous case, we construct a cycle q in $\Gamma(G, X \cup \mathcal{H})$ satisfying (2.15), (2.17) and note that p_1 is an isolated H_λ–component of q. The only additional argument we have to use in this case is that p_1 can not be connected to the H_λ–component of q containing p_2. Indeed, for otherwise we have $g \in H_\lambda$ that contradicts to our assumption. The rest of the proof coincides with that in the previous case. □

In case of torsion free groups we have immediately

Corollary 2.37. *Suppose that G is a torsion free group, G is finitely presented with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$, and the relative Denh function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda})$ is well–defined. Then each subgroup H_λ is malnormal, i.e., $H_\lambda^g \cap H_\lambda = \{1\}$ whenever $g \notin H_\lambda$.*

Let us mention one more corollary of Proposition 2.36.

Corollary 2.38. *Suppose that a group G is finitely presented with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Denh function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda})$ is well–defined. Suppose that $g \in G$ is an element of infinite order and $g^n \in H_\lambda$ for some $n \in \mathbb{Z} \setminus \{0\}$ and some $\lambda \in \Lambda$. Then $g \in H_\lambda$.*

Proof. If $g^n \in H_\lambda$, then the intersection $H_\lambda^g \cup H_\lambda$ contains $\langle g^n\rangle$. Since $\langle g^n\rangle$ is infinite, $g \in H_\lambda$ by Proposition 2.36. □

To formulate our next results, we need an auxiliary notion.

Definition 2.39. A function $f : \mathbb{N} \to \mathbb{N}$ is said to be *subnegative* if

$$f(a+b) \geq f(a) + f(b)$$

for any $a, b \in \mathbb{N}$. Given an arbitrary function $f : \mathbb{N} \to \mathbb{N}$, the *subnegative closure* of f is defined to be

$$\bar{f}(n) = \max_{i=1,\ldots,n} \left(\max_{a_1+\cdots+a_i=n,\ a_i\in\mathbb{N}} (f(a_1) + \cdots + f(a_i)) \right) \tag{2.19}$$

In fact, $\bar{f}$ is the smallest subnegative function such that $\bar{f}(n) \geq f(n)$ for all n.

Below we will speak about subnegative closure of (relative) Dehn functions. It is easy to see that even the ordinary Dehn function of a finitely presented group is not necessarily subnegative. For example, for the group presentation $\langle a \mid a = 1, a^2 = 1\rangle$ we have $\delta(1) = 1$, $\delta(2) = 1$. The question whether or not every Dehn function of a finitely presented group is equivalent to some subnegative function is more complicated. We only note that this question is still open and refer the reader to [47] for more comprehensive discussion.

Theorem 2.40. *Let G be a group, $\{H_1, \ldots H_m\}$, $\{K_1, \ldots, K_l\}$ two finite collections of subgroups of G. Assume that G is finitely presented with respect to $\{H_1, \ldots H_m\} \cup \{K_1, \ldots, K_l\}$ and each subgroup H_i is finitely presented itself. Then G is finitely presented with respect to $\{K_1, \ldots, K_l\}$.*

Moreover, if $\delta_1, \ldots, \delta_m$ *are the ordinary Dehn functions of* $H_1, \ldots, H_m$ *respectively and the relative Dehn function* δ *of* G *with respect to* $\{H_1, \ldots H_m\} \cup \{K_1, \ldots, K_l\}$ *is finite for each value of the argument, then the relative Dehn function* γ *of* G *with respect to* $\{K_1, \ldots, K_l\}$ *is well–defined and satisfies the inequality*

$$\gamma(n) \preceq \bar{f} \circ \delta(n), \tag{2.20}$$

where $\bar{f}$ *is the subnegative closure of the function*

$$f(n) = \max_{i=1,\ldots,m} \delta_i(n).$$

Proof. Before proving the theorem we have to make a few remarks. In the particular case when G is a fundamental group of a complex of groups with finite edge groups, $H_1, \ldots, H_m$ are vertex groups, and $l = 0$ this theorem is equivalent to the main result of the paper [16]. In the case $l = 0$ and the group G is hyperbolic with respect to $\{H_1, \ldots H_m\}$ in the sense of Farb with the BCP property the analogue of Theorem 2.40 can be found in [37]. (However the statement of the theorem in [37] does non involve the subnegative closure, which is required in order to make the proof correct.) Although the proof in the general case exploit similar ideas, we provide it here for convenience of the reader.

By the assumptions of the theorem, the group G is a quotient of the group

$$F = F(X) * \widetilde{H}_1 * \ldots * \widetilde{H}_m * \widetilde{K}_1 * \ldots * \widetilde{K}_l,$$

where $\widetilde{H}_i \cong H_i$ and $\widetilde{K}_j \cong K_j$. Let

$$\mathcal{H} = \bigsqcup_{i=1}^{m} \left(\widetilde{H}_i \setminus \{1\}\right)$$

and

$$\mathcal{K} = \bigsqcup_{j=1}^{l} \left(\widetilde{K}_j \setminus \{1\}\right).$$

We start with a finite reduced relative presentation

$$G = \left\langle X \cup \mathcal{H} \cup \mathcal{K} \;\middle|\; S = 1,\ S \in \bigcup_{i=1}^{m} \mathcal{S}_i,\ P = 1,\ P \in \bigcup_{i=1}^{m} \mathcal{P}_i,\ R = 1,\ R \in \mathcal{R} \right\rangle, \tag{2.21}$$

of the group G with respect to $\{H_1, \ldots H_m\} \cup \{K_1, \ldots, K_l\}$, where $\mathcal{S}_i$ (respectively $\mathcal{P}_i$) is the set of all words over $\widetilde{H}_i$ (respectively $\widetilde{K}_i$) representing the identity in G. Suppose that the groups $H_1, \ldots H_m$ have finite presentations

$$H_i = \langle Y_i \mid T = 1,\ T \in \mathcal{T}_i \rangle, \ \ i = 1, \ldots, m. \tag{2.22}$$

Let O_i denote the set of all letters from $\widetilde{H}_i$ that appear in words $R \in \mathcal{R}$. Since O_i is finite for every i, without loss of generality we can assume that the (finite, symmetrized) generating set Y_i of H_i contains all elements of H_i that are represented by letters from O_i. Thus we can regard words from $\mathcal{R}$ as words in the alphabet

$$Z = X \bigcup \left(\bigcup_{i=1}^{m} Y_i \right) \bigcup \mathcal{K}.$$

We set

$$\mathcal{T} = \bigcup_{i=1}^{m} \mathcal{T}_i.$$

Evidently G can be defined by the finite relative presentation

$$G = \left\langle X \cup \left(\bigcup_{i=1}^{m} Y_i \right) \cup \mathcal{K} \;\middle|\; P = 1,\ P \in \bigcup_{i=1}^{l} \mathcal{P}_i,\ T = 1,\ T \in \mathcal{T},\ R = 1,\ R \in \mathcal{R} \right\rangle \tag{2.23}$$

with respect to $\{K_1, \ldots, K_l\}$. Assume that $\delta_i, i = 1, \ldots m$, are the Dehn function of the presentations (2.22), and δ, γ are the relative Dehn function of the presentations (2.21) and (2.23) respectively.

Let us fix an arbitrary $n \in \mathbb{N}$. Consider a word W over the alphabet Z such that

$$\|W\| \leq n, \tag{2.24}$$

and W represents the identity in G. Since $Z \subseteq X \cup \mathcal{H} \cup \mathcal{K}$, one can regard W as the word over $X \cup \mathcal{H} \cup \mathcal{K}$. We take a van Kampen diagram Δ over (2.21) such that

$$\phi(\partial\Delta) \equiv W \tag{2.25}$$

and assume that Δ has minimal type among all van Kampen diagrams satisfying (2.25). Thus

$$N_{\mathcal{R}}(\Delta) \leq \delta(n). \tag{2.26}$$

Suppose that Π is an S_i–cell in Δ. If $e \in \partial\Pi$ is an internal edge in Δ, then e belongs to the boundary of some $\mathcal{R}$–cell by Lemma 2.15 and therefore $\phi(e) \in Y_i$ according to our choice of Y_i. If $e \in \partial\Pi$ is external, then $\phi(e) \in Y_i$ since W is a word over Z. Thus for any S_i–cell Π of Δ, $\phi(\partial\Pi)$ is a word over Y_i. This observation allows to transform Δ into a van Kampen diagram Θ over (2.23) in the following way. For every S_i–cell Π of Δ, we consider a van Kampen diagram $\Sigma(\Pi)$ over (2.22) whose boundary label coincides with the boundary label of Π and replace Π with $\Sigma(\Pi)$ in Δ. Doing this for all S_i-cells Π in Δ, $i = 1, \ldots, m$, we get a diagram Θ over (2.23).

We want to estimate the number of cells in Θ assuming that for every $\mathcal{S}$–cell Π in Δ, $\Sigma(\Pi)$ has minimal possible number of cells. Let $S(\Delta)$ denote the set of all $\mathcal{S}$–cells of the diagram Δ. We also recall that *Area* $(\Sigma(\Pi))$ denotes the number of all cells in the diagram $\Sigma(\Pi)$.

Using Corollary 2.16, we obtain

$$\sum_{\Pi \in S(\Delta)} l(\partial\Pi) \leq M N_{\mathcal{R}}(\Delta) + l(\partial\Delta), \tag{2.27}$$

where M is the maximum of lengths of the relators from $\mathcal{R}$. Combining (2.24)–(2.27), we have

$$\begin{aligned}\sum_{\Pi\in S(\Delta)} Area\ (\Sigma(\Pi)) &\leq \sum_{\Pi\in S(\Delta)} f(l(\partial\Pi)) \leq \bar{f}\left(\sum_{\Pi\in S(\Delta)} l(\partial\Pi)\right)\\ &\leq \bar{f}(MN_{\mathcal{R}}(\Delta)+\|W\|) \leq \bar{f}(M\delta(n)+n).\end{aligned}$$

Finally, we have the following estimate for the sum of the numbers $N_{\mathcal{R}}(\Theta)$ and $N_{\mathcal{T}}(\Theta)$ of $\mathcal{R}$ and $\mathcal{T}$–cells in Θ respectively:

$$\begin{aligned}N_{\mathcal{R}}(\Theta)+N_{\mathcal{T}}(\Theta) &\leq N_{\mathcal{R}}(\Delta)+\sum_{\Pi\in S(\Delta)} Area\ (\Sigma(\Pi))\\ &\leq \delta(n)+\bar{f}(M\delta(n)+n).\end{aligned}$$

This yields (2.20). □

Corollary 2.41. *Suppose that G is a group hyperbolic relative to a finite collection of finitely presented subgroups $\{H_1,\ldots H_m\}$ and $f(n)$ is an isoperimetric function of H_i for any $i=1,\ldots,m$. Then G is finitely presented itself and $\bar{f}$ is an isoperimetric function of G. In particular, if each of the subgroups H_i is hyperbolic, then G is hyperbolic.*

In the particular case of hyperbolic products of groups, this corollary was obtained by Pankrat'ev [70].

2.4 Splitting Theorem for relatively finitely presented groups

Although the definition of relative Dehn functions has been given in the general situation, by technical reasons it is more convenient to deal with finitely generated groups. In this section we prove the Splitting Theorem for groups given by finite relative presentations, which allows to reduce some questions concerning algebraic properties to the case when the group is finitely generated in the ordinary non–relative sense.

We begin with some basic notions of the Bass–Serre theory of groups acting on trees.

Definition 2.42. A *finite graph of groups* $\mathcal{G}$ consists of the following data.

1) A finite connected oriented graph $\mathcal{G}$; we denote by $E(\mathcal{G})$ and $V(\mathcal{G})$ its set of edges and set of vertices respectively.

2) For every vertex $v\in V(\mathcal{G})$, one associates a group G_v; the groups G_v, $v\in V(\mathcal{G})$, are called *vertex groups*.

3) For every edge $e \in E(\mathcal{G})$, one associates a group G_e together with monomorphisms $\alpha_e : G_e \to G_{e_-}$, $\omega_e : G_e \to G_{e_+}$, where e_- and e_+ are the origin and the terminus of the edge e respectively. The groups G_e, $e \in E(\mathcal{G})$, are called *edge groups.*

Definition 2.43. Let Θ be a maximal tree in $\mathcal{G}$. The fundamental group $\pi_1(\mathcal{G}, \Theta)$ of the finite graph of groups $\mathcal{G}$ at Θ is the group generated by the groups G_v, $v \in V(\mathcal{G})$, and elements t_e, $e \in E(\mathcal{G})$, subject to the relations

$$t_e^{-1}\alpha_e(g)t_e = \omega_e(g), \quad g \in G_e,\ e \in E(\mathcal{G}),$$

$$t_{e^{-1}} = t_e^{-1},\ e \in E(\mathcal{G}),$$

and

$$t_e = 1, \quad \text{if } e \in \Theta.$$

The group $\pi_1(\mathcal{G}, \Theta)$ is independent up to isomorphism of the choice of the maximal tree Θ [77, Prop. 20].

In particular, if $\mathcal{G}$ is a tree itself, then the fundamental group has the presentation

$$\langle G_v,\ v \in V(\mathcal{G}) \mid \alpha_e(g) = \omega_e(g),\ g \in G_e,\ e \in E(\mathcal{G})\rangle . \tag{2.28}$$

Now we formulate the main result of this section.

Theorem 2.44. *Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. Suppose that G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$.*

1) There exist a finite subset $\Lambda_0 = \{\lambda_1, \ldots, \lambda_m\} \subseteq \Lambda$ such that G splits as the free product

$$G = \left(\ast_{\lambda\in\Lambda\setminus\Lambda_0} H_\lambda\right) \ast G_0, \tag{2.29}$$

where G_0 is the subgroup of G generated by $H_{\lambda_1}, \ldots, H_{\lambda_m}$ and X.

2) The groups G_0 is isomorphic to the fundamental group of the tree of groups $\mathcal{G}$ drawn on Fig. 2.7 for some finitely generated groups $L_1, \ldots, L_m$, Q, and the following conditions hold.

a) The group Q is finitely presented with respect to the collection of subgroups $\{L_1, \ldots, L_m\}$.

b) If, in addition, the Denh function δ_G^{rel} of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ is well–defined, then the relative Dehn function δ_Q^{rel} of Q with respect to $\{L_1, \ldots, L_m\}$ is well–defined and

$$\delta_Q^{rel} \preceq \delta_G^{rel} \preceq \bar{\delta}_Q^{rel}, \tag{2.30}$$

where $\bar{\delta}_Q^{rel}$ is the subnegative closure of δ_Q^{rel}



Figure 2.7: The tree of groups $\mathcal{G}$.

Proof. Suppose that G is given by some finite reduced relative presentation (2.3) with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$. We keep our notation from Section 2.2, namely we will use the sets Ω_λ and Ω. For simplicity, we will identify the set Ω_λ, which is a subset of elements of G, with the subset of $\widetilde{H}_\lambda$ consisting of letters representing elements of Ω_λ.

As G is finitely presented relative to $\{H_\lambda\}_{\lambda\in\Lambda}$, there is a finite set of indices $\Lambda_0 = \{\lambda_1, \ldots, \lambda_m\}$ such that no relators from $\mathcal{R}$ involve letters from $\widetilde{H}_\lambda$ for $\lambda \in \Lambda \setminus \Lambda_0$. Evidently we have (2.29), where

$$G_0 = \langle X,\ \mathcal{H}\ | S = 1, S \in \bigcup_{i=1}^{m} \mathcal{S}_{\lambda_i},\ R = 1,\ R \in \mathcal{R}\rangle. \tag{2.31}$$

We divide the rest of the proof into a few lemmas. For every $i = 1, \ldots, m$, we consider the group $L_i = \langle \Omega_i \rangle$. Let

$$F_Q = F(X) * \widetilde{L}_1 * \ldots * \widetilde{L}_m,$$

where $\widetilde{L}_i$ are isomorphic copies od L_i. Set

$$\mathcal{L} = \bigsqcup_{i=1}^{m} \left(\widetilde{L}_i \setminus \{1\}\right).$$

By $\mathcal{T}_i$ we denote the set of all words over the alphabet $\widetilde{L}_i \setminus \{1\}$ representing the identity in the group G. Let

$$\mathcal{T} = \bigcup_{i=1}^{m} \mathcal{T}_{\lambda_i}.$$

Lemma 2.45. *Let Q be the subgroup of G generated by the set $\Omega \cup X$. Then Q has the finite relative presentation*

$$\langle X, \mathcal{L} \mid T = 1,\ T \in \mathcal{T},\ R = 1,\ R \in \mathcal{R}\rangle. \tag{2.32}$$

with respect to $\{L_1, \ldots, L_m\}$. Moreover, the relative Dehn function of (2.32) satisfies the inequality

$$\delta_Q^{rel} \preceq \delta_G^{rel}.$$

Proof. Indeed, let W be a word over $X \cup \mathcal{L}$ of length n that represents 1 in the group G. To prove the lemma we have to show that there exists a van Kampen diagram over (2.32) with boundary label W and number of cells at most $\delta_G^{rel}(n)$.

To this end we note that any word over $X \cup \mathcal{L}$ can be regarded as a word over $X \cup \mathcal{H}$. We consider a diagram Θ of minimal type over (2.3) such that $\phi(\partial\Delta) \equiv W$. Suppose that e is an edge of Θ labelled by a letter from $\mathcal{H}$. If e is internal, e belongs to the boundary of some $\mathcal{R}$–cell by Lemma 2.15. Hence $\phi(e)$ represents an element w of $\Omega_{\lambda_i} \subseteq L_i$ for some i and we can regard $\phi(e)$ as an element of $\widetilde{L}_i \setminus \{1\}$. Thus if Π is an S–cell in Θ, then $\partial\Pi$ is labelled by a word in the alphabet $\widetilde{L}_i \setminus \{1\}$ for a certain $i \in \{1, \ldots, m\}$. As $\mathcal{T}_i$ contains all words over $\widetilde{L}_i \setminus \{1\}$ representing the identity in G, one can think of Π as a cell corresponding to a relator from $\mathcal{T}_i$. Thus Θ can be regarded as a diagram over (2.32). This completes the proof. □

Let $\alpha_i : L_i \to Q$ and $\omega_i : L_i \to H_{\lambda_i}$ be the natural embeddings. Then the fundamental group of the graph of groups $\mathcal{G}$ can be represented as

$$\langle Q, H_{\lambda_i}, i = 1, \ldots, m \mid \alpha_i(l) = \omega_i(l), l \in L_i, i = 1, \ldots, m \rangle . \tag{2.33}$$

It is easy to see that using (2.32), we can obtain (2.31) from (2.33) by applying Tietze transformations. This proves that G_0 splits at the fundamental group of the graph of groups $\mathcal{G}$.

Lemma 2.46. *We have $\delta_G^{rel} \preceq \bar{\delta}_Q^{rel}$.*

Proof. Let q be a cycle in $\Gamma(G, X \cup \mathcal{H})$ of length $l(q) \leq n$. We repeat here the trick used in the proof of Lemma 2.27. Let $q = q_1 r_1 \ldots q_n r_n$ be the decomposition of q into the product of components $q_1, \ldots, q_n$ and subpath $r_1, \ldots, r_n$ labelled by words in X (r_i may be trivial for some i). By $\mathcal{Q}$ we denote the set of all irreducible cycles q' in $\Gamma(G, X \cup \mathcal{H})$ with the same initial point as q and such that q' can be represented as

$$q' = q'_1 r_1 \ldots q'_n r_n, \tag{2.34}$$

where q'_i is an H_μ–component of q' for the same μ as q_i and $(q'_i)_- = (q_i)_-$, $(q'_i)_+ = (q_i)_+$ for $i = 1, 2, \ldots, n$. Let $\mathcal{D}$ be the set of all diagrams over (2.3) with boundary label $\phi(q')$ for some $q' \in \mathcal{Q}$.

Take the diagram Δ of minimal type in $\mathcal{D}$. Then Δ can be represented as the union of subdiagrams $\Delta_1, \Delta_2, \ldots, \Delta_k$, each Δ_i is homeomorphic to a disk, connected by some paths entirely consisting of external edges of the second type (see Fig. 2.8).

Repeating the arguments from the proof of Lemma 2.27 and using our assumptions about Δ, one can easily show that no $\mathcal{S}$–cell of Δ has a common edge



Figure 2.8: The decomposition of Δ.

with $\partial\Delta_i$ for some i. Therefore, for any i, every edge e of Δ_i belongs to the boundary of some $\mathcal{R}$–cell. Hence $\phi(e)$ can be regarded as a letter from $X \cup \mathcal{L}$. As $\mathcal{T}_i$ contains all words over $\widetilde{L}_i \setminus \{1\}$ representing the identity in G, this allows us to consider Δ_i, $i = 1, \ldots, k$, as diagrams over (2.32). We obtain

$$Area^{rel}(q) = \sum_{i=1}^{k} N_{\mathcal{R}}(\Delta_i) \leq \sum_{i=1}^{k} \delta_Q^{rel}(l(\partial\Delta_i)) \leq \bar{\delta}_Q^{rel}\left(\sum_{i=1}^{k} l(\partial\Delta_i)\right) \leq \bar{\delta}_Q^{rel}(n).$$

□

□

We notice one corollary of Theorem 2.44.

Corollary 2.47. *If the group G in the Theorem 2.44 is hyperbolic with respect to the collection $\{H_\lambda\}_{\lambda\in\Lambda}$, then Q is hyperbolic with respect to the collection $\{L_1, \ldots, L_m\}$.*

The Splitting Theorem together with the Gruško–Neumann theorem [56, Proposition 3.7] also imply

Corollary 2.48. *Let G be a finitely generated group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of nontrivial subgroups of G. Suppose that G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$. Then the collection $\{H_\lambda\}_{\lambda\in\Lambda}$ is finite, i.e., card $\Lambda < \infty$.*

Proof. Given a finitely generated group K, by $rank\, K$ we denote the minimal number of generators of K. By the Gruško–Neumann theorem, we have

$$rank\, G_0 + \sum_{\lambda\in\Lambda\setminus\Lambda_0} rank\, H_\lambda \leq rank\, G < \infty. \tag{2.35}$$

As Λ_0 is finite, the corollary follows immediately from (2.35). $\square$

We conclude this section with one more result describing the geometry of the embedding $Q \to G$. This result will be used in Section 4.3 (as well as the Splitting Theorem) to study cyclic subgroups of relatively hyperbolic group. Besides it seems to be of independent interest.

Proposition 2.49. *Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. Suppose that G is finitely presented with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$. Let Q and L_i, $i = 1, \ldots, m$ be the groups provided by Theorem 2.44. We denote by $dist_{X\cup\mathcal{L}}$ the relative metric on Q with respect to the collection $\{L_1, \ldots, L_m\}$. Then the mapping of metric spaces $(Q, dist_{X\cup\mathcal{L}}) \to (G, dist_{X\cup\mathcal{H}})$ induced by the natural embedding $Q \to G$ is an isometry.*

Proof. It is easy to see that $|g|_{X\cup\mathcal{L}} \geq |g|_{X\cup\mathcal{H}}$ for any $g \in G$. Let us prove the converse inequality.

Suppose that $g \in Q$ is an element such that the relative length of g in Q satisfies $|g|_{X\cup\mathcal{L}} > |g|_{X\cup\mathcal{H}}$. Let W_1 and W_2 be shortest words over $X \cup \mathcal{L}$ and $X \cup \mathcal{H}$ respectively representing the element g. In what follows we regard $\mathcal{L}$ as a subset of $\mathcal{H}$.

Consider the cycle pq^{-1} in $\Gamma(G, X \cup \mathcal{H})$ such that

$$\phi(p) \equiv W_1, \qquad \phi(q) \equiv W_2.$$

Let s be an H_λ–component of q^{-1} for some λ. We are going to show that $\lambda \in \Lambda_0$ and $\overline{\phi(s)} \in L_i$ for some $i \in \{1, \ldots, m\}$.

If s is an isolated H_λ–component of pq^{-1}, $\overline{\phi(s)} \in \langle \Omega_\lambda \rangle$ by Lemma 2.27. Note that Ω_λ is non–empty only if $\lambda \in \Lambda_0$. Thus $\overline{\phi(s)} \in L_i$ for some $i \in \{1, \ldots, m\}$. Further suppose that s is not isolated in pq^{-1}. Let $t_1, \ldots, t_k, s$ be the maximal connected system of H_λ–components of pq^{-1} containing s. Since the word W_2 is a shortest words over $X \cup \mathcal{H}$ representing the element g, the path q is geodesic in $\Gamma(G, X \cup \mathcal{H})$. Hence any component of q is isolated in q. It follows that $t_1, \ldots, t_k$ are subpaths of p. Therefore, $\phi(t_1), \ldots, \phi(t_k)$ are H_λ–syllables in W_1. Thus $\lambda = \lambda_i$ for some i and $\overline{\phi(t_j)} \in L_i$ for $j = 1, \ldots, k$. Finally, using Corollary 2.28, we obtain

$$\overline{\phi(s)} \in \langle \Omega_{\lambda_i}, \overline{\phi(t_1)}, \ldots, \overline{\phi(t_k)} \rangle = L_i.$$

We have proved that any H_λ–syllable of W_2 represents an element of L_i for some i. This means that one can think of W_2 an a word over $X \cup \mathcal{L}$. However,

$$\|W_2\| = |g|_{X\cup\mathcal{L}} < |g|_{X\cup\mathcal{H}} = \|W_1\|$$

that contradicts to the minimality of W_1. $\square$

2.5 Isoperimetric functions of Cayley graphs.

In order to apply various well–known results about hyperbolic spaces to a pair $(G, \{H_\lambda\}_{\lambda\in\Lambda})$ with linear relative Dehn function, we have to study more extensively the geometry of the corresponding Cayley graph $\Gamma(G, X\cup\mathcal{H})$. In the present section we establish the relation between the linearity of the Dehn function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ and hyperbolicity of $\Gamma(G, X\cup\mathcal{H})$.

We begin with a definition of the notion of area of a cycle in a metric graph, which is a particular case of a more general concept (see [46, Sec. 5.F], [19, Ch. III.H, Sec.2] or [1], [6] for a refined version of area).

First we have to introduce an auxiliary terminology. Let D^n denote the unit disk in the n–dimensional Euclidean space. Recall that a *combinatorial map* between CW complexes is a map that sends open cells homeomorphically onto open cells, and a *combinatorial complex* is a CW complex K such that for any $(n+1)$–cell e of K, the attaching map $\phi_e : \partial D^{n+1}\to Sk^{(n)}K$, where $Sk^{(n)}K$ denotes the n–skeleton of K, is combinatorial with respect to some combinatorial structure on $\partial D^{n+1}=\mathbb{S}^n$. (This definition involves the induction on dimension.) In what follows we will also work with a lager category of maps. One defines a *singular combinatorial map* $f : L\to K$ between CW complexes to be a continuous map such that for every open n–cell e in L, either $f|_e$ is a homeomorphism onto an open cell of K or else $f(e)$ is contained in the $(n-1)$–skeleton of L [18].

Definition 2.50. (k–filling, $A^{(k)}$, and Isoperimetric Inequality). Let $k\geq 3$. A *k–partition* of D^2 is a homeomorphism P from D^2 to a combinatorial 2–complex in which every 2–cell is an l–gone for some $3\leq l\leq k$. We endow D^2 with the induced cell structure and refer to the preimages under P of 0–cells, 1–cells, and 2–cells as, respectively, vertices, edges, and faces of D^2. We also denote by $Sk^{(1)}D^2$ the 1–skeleton of D^2 with the induced cell structure.

Let Σ be a graph equipped with the combinatorial metric, that is, each edge of Σ has length 1. Let c be a combinatorial cycle in Σ. A *k–filling* of c consists of a k–partition P of D^2 and a singular combinatorial map $\Phi : Sk^{(1)}D^2\to\Sigma$ such that $\Phi(\partial D^2)=c$. In this situation we write $|\Phi|$ to denote the number of faces of D^2 with respect to the cell structure induced by P. The *k–area* of c is defined to be

$$A^{(k)}(c)=\min\{|\Phi|,\ \Phi \text{ is a } k\text{−filling of c}\}$$

(if there is no k–filling of c, we put $A^{(k)}(c)=\infty$).

We also set

$$f^{(k)}_\Sigma(n)=\sup_{l(c)\leq n} A^{(k)}(c),$$

where the supremum is taken over all cycles of combinatorial length at most n.

It is easy to check that if for some $k\geq 3$ any cycle in Σ admits a k–filling and the corresponding function $f^{(k)}_\Sigma$ is finite for each value of the argument, then for any $k' \geq k$, $f_{\Sigma}^{(k')}$ is finite for each value of the argument and, moreover, $f_{\Sigma}^{(k')} \sim f_{\Sigma}^{(k)}$. (The proof of this fact in a much more general situation can be found in [19, Ch. III.H, Sec.2].) In this case $f_{\Sigma}^{(k)}$ (regarded up to equivalence) is called the *Dehn function* of Σ and is denoted by f_{Σ}. As above, we consider Dehn functions of metric graphs up to the usual equivalence.

Example 2.51. In case Σ is the Cayley graph of a group Q given by a finite presentation $\langle Y \mid \mathcal{Q}\rangle$, any cycle in Σ admits a k–filling for $k = \max\limits_{Q\in\mathcal{Q}} \|Q\|$ and $f^{(k)}(n) < \infty$ for any n. Moreover, $f^{(k)}$ is equivalent to the ordinary Dehn function of G. The assumption of finite presentability of G is essential, since if G is not finitely presented, then for any $k \in \mathbb{N}$, there exists a cycle c is Σ that admits no k–filling.

If we consider a group G which is finitely presented with respect to a collection of the subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$, then the relation between the relative Dehn function δ_G^{rel} of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ and the Dehn function $f_{\Gamma(G,X\cup\mathcal{H})}$ of $\Gamma(G, X\cup\mathcal{H})$ can be expressed as follows.

Lemma 2.52. *Suppose G is a group with a reduced finite relative presentation (2.3) with respect to a collection of the subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$. Set*

$$k = \max\{\max_{R\in\mathcal{R}} \|R\|, 4\}. \tag{2.36}$$

Then every cycle q in $\Gamma(G, X\cup\mathcal{H})$ admits a k–filling and we have

$$A^{(k)}(q) \leq (k+1)Area^{rel}(q) + 2l(q).$$

Proof. We consider a van Kampen diagram Δ of minimal type over (2.3) such that $\phi(\partial\Delta) \equiv \phi(q)$. In particular, this means that

$$N_{\mathcal{R}}(\Delta) = Area^{rel}(q). \tag{2.37}$$

Using Corollary 2.16, we obtain

$$\sum_{\Pi\in S(\Delta)} l(\partial\Pi) \leq \max_{R\in\mathcal{R}} \|R\| Area^{rel}(q) + l(q) \leq kArea^{rel}(q) + l(q), \tag{2.38}$$

where $S(\Delta)$ is the set of all $\mathcal{S}$–cells of Δ. Now we obtain a diagram Ψ from Δ as follows. Let us take a cell $\Pi \in S(\Delta)$. If $l(\partial\Pi) > 3$, we can "triangulate" Π by considering a van Kampen diagram $\Upsilon(\Pi)$ over (2.3) with the same boundary label as Π such that every cell of $\Upsilon(\Pi)$ is an $\mathcal{S}$–cell whose boundary has length 3. Obviously the minimal number of cells in such a diagram satisfies

$$Area(\Upsilon(\Pi)) \leq l(\partial\Pi) - 2. \tag{2.39}$$

Then we replace the cell Π of Δ with the diagram $\Upsilon(\Pi)$. Doing this for all $\Pi \in S(\Delta)$, we obtain a new diagram Ψ over (2.3) in which the boundary of every S–cell has length at most 3. Combining (2.37), (2.38), and (2.39), we obtain the following estimate on the number of cells in Ψ:

$$\begin{aligned} Area(\Psi) &= N_{\mathcal{R}}(\Delta) + \sum_{\Pi \in S(\Delta)} Area(\Upsilon(\Pi)) \\ &\leq Area^{rel}(q) + \sum_{\Pi \in S(\Delta)} l(\partial \Pi) \leq (k+1) Area^{rel}(q) + l(q). \end{aligned}$$

We are going to define a k–filling of q using Ψ. Obviously every 2–cell of Ψ is an l–gone for some $l \leq k$. The only difficulty is that Ψ may not be homeomorphic to a disk. However in this case we can transform Ψ into a simply–connected diagram Ψ' by using the so-called 0–bordering of the contour of Ψ, which is a particular case of a 0-refinement (we refer the reader to [61, Ch.4, Section 11.5] for the definition). Applying this process, we add new cells to Ψ, the so–called 0–faces, such that the contour of every 0–face has length 4 and the image of every 0–face under the canonical map into the Cayley graph $\Gamma(G, X \cup \mathcal{H})$ is an edge (this is one of the reasons why we need singular maps in the definition of the k–filling). Moreover, the number of additional cells equals $l(q)$. The diagram Ψ' defines a k–partition of D^2 and the natural map $Sk^{(1)}\Psi' \to \Gamma$ such that $\partial \Psi'$ is mapped onto q gives a k–filling of q. Thus we have

$$A^{(k)}(q) \leq Area(\Psi') = Area(\Psi) + l(q) \leq (k+1) Area^{rel}(q) + 2l(q).$$

□

From the above lemma we obtain

Theorem 2.53. *Suppose that a group G is finitely presented with respect to a collection of subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$ and the Dehn relative function δ_G^{rel} of G with respect to $\{H_\lambda\}_{\lambda \in \Lambda}$ is well–defined. Then the Denh function $f_{\Gamma(G, X \cup \mathcal{H})}$ of the Cayley graph $\Gamma(G, X \cup \mathcal{H})$ is equivalent to δ_G^{rel}.*

Proof. Lemma 2.52 gives us the estimate

$$f_{\Gamma(G, X \cup \mathcal{H})} \preceq \delta_G^{rel}.$$

To prove the reverse inequality, we take k such that any cycle in $\Gamma(G, X \cup \mathcal{H})$ admits a k–filling. Let q be such a cycle, $l(q) \leq n$. Suppose that $\Phi : Sk^{(1)} D^2 \to \Gamma(G, X \cup \mathcal{H})$ is a k–filling having the minimal number of faces among all k–fillings of q. Thus we have

$$|\Phi| \leq f_{\Gamma(G, X \cup \mathcal{H})}(n). \tag{2.40}$$

Using Φ we can define labels and orientations on edges of D^2 as follows. If e is an edge of D^2 and $\Phi(e)$ is an edge of $\Gamma(G, X \cup \mathcal{H})$, we endow e with the induced label and orientation in the obvious way; if $\Phi(e)$ is a vertex of $\Gamma(G, X \cup \mathcal{H})$, then we put $\phi(e) = 1$ (the orientation does not matter in this case). Further, let Π

be a cell of D^2. Since $\Phi(\partial\Pi)$ has length at most k, there exists a van Kampen diagram $\Xi(\Pi)$ over (2.3) such that $\phi(\partial\Xi(\Pi)) \equiv \phi(\partial\Pi)$ and

$$N_{\mathcal{R}}(\Xi(\Pi)) \leq \delta_G^{rel}(k) < \infty. \tag{2.41}$$

Finally we can obtain a (0–refinement of) van Kampen diagram over (2.3) such that $\phi(\partial\Delta) \equiv \phi(q)$ by replacing all cells Π of D^2 with the corresponding diagrams $\Xi(\Pi)$. Evidently, from (2.40) and (2.41) we obtain

$$Area^{rel}(q) \leq N_{\mathcal{R}}(\Delta) \leq |\Phi|\delta_G^{rel}(k) \leq f_{\Gamma(G,X\cup\mathcal{H})}(n)\delta_G^{rel}(k).$$

This leads to the inequality $\delta_G^{rel} \preceq f_{\Gamma(G,X\cup\mathcal{H})}$. ☐

Corollary 2.54. *Suppose that a group G is finitely presented with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Dehn function δ_G^{rel} of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$) is well–defined. Then the following conditions are equivalent.*

1) G is hyperbolic relative to $\{H_\lambda\}_{\lambda\in\Lambda}$.

2) The Cayley graph $\Gamma(G, X\cup\mathcal{H})$ is a hyperbolic metric space (we refer to Section 3.1 for the definition).

Proof. By Theorem 2.53 the relative Dehn function of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ is linear if and only if the Dehn function of $\Gamma(G, X\cup\mathcal{H})$ is linear. As is well–known, the las condition is equivalent to the hyperbolicity of $\Gamma(G, X\cup\mathcal{H})$. Up to notation, the proof can be found in [19, Ch. III.H, Sec. 2]; see also [57], [62], [79]. ☐

Example 2.55. The requirement $\delta_G^{rel}(n) < \infty$ for any n in the statement of Corollary 2.54 is essential. Indeed, consider the group $G \cong \mathbb{Z}$ and a subgroup H of finite index in G. Then obviously $\Gamma(G, X\cup\mathcal{H})$ has finite diameter. In particular, $\Gamma(G, X\cup\mathcal{H})$ is hyperbolic. However, δ_G^{rel} is not well–defined by Proposition 2.36.

Definition 2.56. A function $f : \mathbb{N} \to \mathbb{N}$ is said to be *subquadratic* if $f = o(n^2)$ as $n \to \infty$.

Recall that every finitely presented group with subquadratic Dehn function is hyperbolic. This insight is due to M. Gromov [45], and was clarified by Ol'shanskii [62] and others (see [10], [73] and references therein). Our next goal is to show that if G has a subquadratic relative Dehn function δ_G^{rel} with respect to a collection $\{H_\lambda\}_{\lambda\in\Lambda}$, then , in fact, δ_G^{rel} is linear and $\Gamma(G, X\cup\mathcal{H})$ is a hyperbolic metric space.

Corollary 2.57. *Suppose that a group G is finitely presented with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$ and the relative Denh function δ_G^{rel} of G with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$), is well–defined. Then the following conditions are equivalent.*

1) δ_G^{rel} is subquadratic.

2) δ_G^{rel} is linear.

Proof. If δ_G^{rel} is subquadratic, then the Dehn function of $\Gamma(G, X \cup \mathcal{H})$ is equivalent to δ_G^{rel} by Theorem 2.53. It is well–known that if a graph possess a subquadratic Dehn function, it possess a linear one (see [19, Ch. III.H, Sec. 2], [10]). Applying Theorem 2.53 again, we obtain that δ_G^{rel} is linear. $\square$

Chapter 3

Geometry of finitely generated relatively hyperbolic groups

3.1 Conventions and notation

In this chapter we study the geometry of the Caley graph of a finitely generated in the usual non–relative sense group G which is hyperbolic relative to a collection $H_1, \ldots, H_m$ of subgroups. (Recall that by Corollary 2.48, if a finitely generated group is hyperbolic relative to a collection of subgroups, then the collection is finite). Throughout the next three sections we accept the following technical agreements about G and $H_1, \ldots, H_m$.

(i) *G is represented by a finite relative presentation*

$$G = \langle X, H_1, \ldots, H_m \mid R = 1, R \in \mathcal{R} \rangle \tag{3.1}$$

with respect to $\{H_1, \ldots, H_m\}$ and the relative Dehn function of (3.1) satisfies

$$\delta_G^{rel}(n) \leq Ln \tag{3.2}$$

for some constant L; also, as in the previous chapter, we set

$$M = \max_{R \in \mathcal{R}} ||R||.$$

By technical reasons it is convenient to increase L in order to satisfy the inequality

$$ML > 1.$$

(ii) *G is generated by the set X in the ordinary sense.*

(iii) *The set X is chosen in such a way that the following lemma holds.*

Lemma 3.1. *Let G be a finitely generated group hyperbolic relative to a collection of subgroups $\{H_1, \ldots, H_m\}$. Then there exists a finite generating set X of G satisfying the following condition. Let q be a cycle in $\Gamma(G, X \cup \mathcal{H})$, $p_1, \ldots, p_k$ a certain set of isolated H_i–components of q. Then*

$$\sum_{i=1}^{k} dist_X((p_i)_-, (p_i)_+) \leq MLl(q). \tag{3.3}$$

Let us show that we can always ensure the fulfillment of (i)–(iii). We start with an arbitrary finite reduced relative presentation

$$G = \langle X', H_1, \ldots, H_m \mid R = 1, R \in \mathcal{R}' \rangle. \tag{3.4}$$

Let Ω be the set given by Definition 2.25. Note that Ω is finite in our case. We set

$$X = X' \cup \Omega$$

and

$$\mathcal{R} = \mathcal{R}' \cup \{\omega W_\omega^{-1},\ \omega \in \Omega\},$$

where W_ω is a fixed word over X' representing the element ω in G. It is clear that for X and $\mathcal{R}$ chosen in this way, the presentation (3.1) can be obtained from (3.4) by a finite number of Tietze transformations. It remains to prove Lemma 3.1.

Let u_i (respectively v_i) denote $(p_i)_-$ (respectively $(p_i)_+$) regarded as an element of G. By Lemma 2.27, we have

$$\sum_{i=1}^{k} |u_i^{-1} v_i|_\Omega \leq M'L'l(q),$$

where

$$M = \max_{R \in \mathcal{R}'} ||R|| \leq M,$$

and the relative Dehn function of the presentation (3.4) does not exceed $L'n$. Increasing the constant L in (3.2) if necessary, we may assume that $L' \leq L$. Thus we obtain

$$\sum_{i=1}^{k} dist_X((p_i)_-, (p_i)_+) \leq \sum_{i=1}^{k} |u_i^{-1} v_i|_X \leq \sum_{i=1}^{k} |u_i^{-1} v_i|_\Omega \leq M'L'l(q) \leq MLl(q).$$

Definition 3.2. Recall that a metric space Y is called *δ–hyperbolic* (or simply *hyperbolic*) if it satisfies the following *Rips condition.* For any geodesic triangle, each side of the triangle belongs to the union of the closed δ–neighborhood of the other two sides.

By Corollary 2.54, the Cayley graph $\Gamma(G, X \cup \mathcal{H})$ is hyperbolic. In what follows we denote by δ the hyperbolicity constant of $\Gamma(G, X \cup \mathcal{H})$.

3.2 Properties of quasi–geodesics

First of all we recall some facts concerning quasi–geodesics in hyperbolic metric spaces. Below we assume all paths under consideration to be rectifiable (i.e., to have finite length).

Definition 3.3. A path q in a metric space Y is said to be (λ, c)*–quasi–geodesic* for some $\lambda \geq 1$, $c \geq 0$, if for every subpath q of p the inequality

$$l(q) \leq \lambda \, dist(q_-, q_+) + c$$

holds.

The following useful lemma is quite obvious.

Lemma 3.4. *Let p be a (λ, c)–quasi–geodesic in a metric space Y, e a path of length k such that $e_- = p_+$. Then the path pe is $(\lambda, c+(\lambda+1)k)$–quasi–geodesic.*

Proof. Let q be a subpath of pe, q_0 the maximal common subpath of q and p. Then

$$\begin{aligned} l(q) &\leq l(q_0) + k \leq \lambda dist_{X\cup\mathcal{H}}((q_0)_-, (q_0)_+) + c + k \\ &\leq \lambda(dist_{X\cup\mathcal{H}}((q)_-, (q)_+) + k) + c + k. \end{aligned}$$

□

Next result is well known and can be found in [45] or [44].

Lemma 3.5. *For any $\delta \geq 0$, $\lambda \geq 1$, $c \geq 0$, there exists a constant $H = H(\delta, \lambda, c)$ with the following property. If Y is a δ–hyperbolic space and p, q are (λ, c)–quasi–geodesic paths in Y with same endpoints, then p and q belong to the closed H–neighborhoods of each other.*

From the definition of a hyperbolic space, one obtains

Lemma 3.6. *Let Y be a δ–hyperbolic metric space, $Q = p_1p_2p_3p_4$ a geodesic quadrangle in Y. Then each side of Q belongs to the closed 2δ–neighborhood of the other three sides.*

Corollary 3.7. *Let Y be a δ–hyperbolic space and p, q are geodesic paths in Y such that $dist(p_-, q_-) \leq k$ and $dist(p_+, q_+) \leq k$, then p and q belong to the closed $(k + 2\delta)$–neighborhood of each other.*

From the above corollary and Lemma 3.5, we can easily derive

Lemma 3.8. *For any $\delta \geq 0$, $\lambda \geq 1$, $c \geq 0$, and $k \geq 0$, there exists a constant $K = K(\delta, \lambda, c, k)$ such that the following condition holds. Suppose that Y is a δ–hyperbolic space and p, q are (λ, c)–quasi–geodesic paths in Y such that $dist(p_-, q_-) \leq k$ and $dist(p_+, q_+) \leq k$; then p and q belong to the closed K–neighborhoods of each other.*

Given the data described in Section 3.1, there are two Cayley graphs, namely $\Gamma(G,X)$, the Cayley graph of G with respect to the generating set X, and $\Gamma(G, X\cup\mathcal{H})$, the Cayley graph of G with respect to the generating set $X\cup\mathcal{H}$. Obviously we have a natural embedding

$$\iota : \Gamma(G,X) \to \Gamma(G, X\cup\mathcal{H}),$$

which is bijective on the set of vertices. For simplicity we will identify $\Gamma(G,X)$ with its image in $\Gamma(G, X\cup\mathcal{H})$ under the embedding ι.

Assuming the length of each edge of $\Gamma(G, X\cup\mathcal{H})$ to be equal to 1, we obtain combinatorial metrics $dist_X$ and $dist_{X\cup\mathcal{H}}$ in $\Gamma(G,X)$ and $\Gamma(G, X\cup\mathcal{H})$ respectively. When speaking about geodesics (quasi–geodesics) in $\Gamma(G,X)$ or in $\Gamma(G, X\cup\mathcal{H})$ we always mean geodesics (quasi–geodesics) with respect to corresponding combinatorial metric. Note that the restrictions of $dist_X$ and $dist_{X\cup\mathcal{H}}$ on the vertex sets $V(\Gamma(G,X) = V(\Gamma(G,X\cup\mathcal{H})) = G$ coincide with the word metric on G with respect to the generating sets X and $X\cup\mathcal{H}$.

Definition 3.9. A path p in $\Gamma(G, X\cup\mathcal{H})$ is called a *path without backtracking* if for any $i = 1,\ldots,m$, every H_i–component of p is isolated.

Definition 3.10. Let p be a path in $\Gamma(G, X\cup\mathcal{H})$, v a vertex of a component s of p. If $v \neq s_-$ and $v \neq s_+$, we say that v is an inner vertex of s. A vertex u of p is called *non–phase*, if u is a inner vertex of some component of p. All other vertices of p are called *phase*.

Definition 3.11. A path p in $\Gamma(G, X\cup\mathcal{H})$ is said to be *locally minimal* if for any $i = 1,\ldots,m$, every H_i–component of p has length 1 or, equivalently, every vertex of p is phase.

We need the following simple observation.

Lemma 3.12. *Let p be a path in $\Gamma(G, X\cup\mathcal{H})$. Then there exists a locally minimal path $\hat{p}$ having the same set of phase vertices as p. Moreover, if p is (λ, c)–quasi–geodesic, then $\hat{p}$ is (λ, c)–quasi–geodesic; if p is a path without backtracking, then so is $\hat{p}$.*

Proof. Given an H_i–component s of p, we can replace s with the single edge of $\Gamma(G, X\cup\mathcal{H})$ having the label $\phi(e) =_{H_i} \phi(s)$. Doing this for all components of p we obtain the desirable path $\hat{p}$. The verification of the additional properties is straightforward. $\square$

Definition 3.13. Let p, q be two paths in $\Gamma(G, X\cup\mathcal{H})$. We say that p and q are *k–similar* if

$$\max\{dist_X(p_-, q_-),\ dist_X(p_+, q_+)\} \leq k. \tag{3.5}$$

The next proposition is an improved version of Lemma 3.8 for relatively hyperbolic groups.

Definition 3.14. We say that two paths p and q in $\Gamma(G, X \cup \mathcal{H})$ are k–similar if

$$\max\{dist_X(p_-, q_-),\ dist_X(p_+, q_+)\} \leq k.$$

Proposition 3.15. *For any $\lambda \geq 1$, $c \geq 0$, $k \geq 0$ there exists a constant $\varepsilon = \varepsilon(\lambda, c, k) > 0$ having the following property. Let p and q be two k–similar (λ, c)–quasi–geodesic paths in $\Gamma(G, X \cup \mathcal{H})$ (with respect to the relative metric $dist_{X\cup\mathcal{H}}$) such that p is a path without backtracking. Then for any phase vertex u of p there exists a phase vertex v of q such that*

$$dist_X(u, v) \leq \varepsilon. \tag{3.6}$$

Proof. Before proving the theorem, we note that (3.6) is much stronger that the inequality $dist_{X\cup\mathcal{H}}(u, v) \leq \varepsilon$, which follows from Lemma 3.8. In view of Lemma 3.12, it is sufficient to prove the proposition for locally minimal quasi–geodesic paths. Thus we assume that every vertex of p and q is phase.

Let u be a (phase) vertex of p. Recall that $\Gamma(G, X \cup \mathcal{H})$ is a δ–hyperbolic space. We set $K_0 = K(\delta, \lambda, c, k)$ and $K = K(\delta, \lambda, c, K_0) + 1/2$, where $K(\delta, \lambda, c, k)$ and $K(\delta, \lambda, c, K_0)$ are constants provided by Lemma 3.8. Without loss of generality we may assume that $K \in \mathbb{N}$ and

$$K \geq K_0 \geq k. \tag{3.7}$$

Let us choose vertices u_1, u_2 on p as follows. If $dist_{X\cup\mathcal{H}}(p_-, u) \leq 2K$ (respectively $dist_{X\cup\mathcal{H}}(u, p_+) \leq 2K$), we put $u_1 = p_-$ (respectively $u_2 = p_+$). If $dist_{X\cup\mathcal{H}}(p_-, u) > 2K$ (respectively $dist_{X\cup\mathcal{H}}(u, p_+) > 2K$), we take u_1 on the segment $[p_-, u]$ of the path p (respectively u_2 on the segment $[u, p_+]$ of the path p) such that

$$dist_{X\cup\mathcal{H}}(u, u_i) = 2K \tag{3.8}$$

for $i = 1$ (respectively $i = 2$).

Further, by Lemma 3.8, there exist two points v_1, v_2 of q such that

$$dist_{X\cup\mathcal{H}}(v_i, u_i) \leq K_0, \quad i = 1, 2. \tag{3.9}$$

Without loss of generality we may assume that v_1, v_2 are vertices of q. Moreover, we assume that in case $u_1 = p_-$ (respectively $u_2 = p_+$) the vertex v_1 coincides with q_- (respectively the vertex v_2 coincides with q_+).

We denote by p_1, p_2 (respectively by q_0) the segments of p (respectively the segment of q) such that $(p_1)_- = u_1$, $(p_1)_+ = u$, $(p_2)_- = u$, $(p_2)_+ = u_2$ (respectively $(q_0)_- = v_1$, $(q_0)_+ = v_2$). We also denote by o_1, o_2 the paths in $\Gamma(G, X \cup \mathcal{H})$ such that $(o_i)_- = u_i$, $(o_i)_+ = v_i$, $i = 1, 2$ (see Fig. 3.8) and o_1, o_2 are chosen according to the following agreement. If $u_1 = p_-$ (respectively $u_2 = p_+$), then o_1 (respectively o_2) is a geodesic path in $\Gamma(G, X)$. If $u_1 \neq p_1$ (respectively $u_2 \neq p_+$), then o_1 (respectively o_2) is geodesic in $\Gamma(G, X \cup \mathcal{H})$. It follows from (3.5) and (3.7) in the first case and from (3.9) in the second case that

$$l(o_i) \leq K_0. \tag{3.10}$$



Figure 3.1:

Let V denote the set of all vertices z on q_0 that are closest to u, i.e., satisfy the condition

$$dist_{X\cup\mathcal{H}}(u,z)=\min_{v\in q_0} dist_{X\cup\mathcal{H}}(u,v),$$

where v ranges among all vertices of q_0. Taking into account (3.9) and Lemma 3.8 we obtain

$$dist_{X\cup\mathcal{H}}(u,z)\le K(\delta,\lambda,c,K_0)+1/2=K \tag{3.11}$$

for any $z\in V$. To each $z\in V$, we associate the set $O(z)$ of all a geodesic paths o such that $o_-=u$, $o_+=z$. Each of the paths $o\in O(z)$ cuts the cycle $p_1p_2o_2q_0^{-1}o_1^{-1}$ into two parts denoted c_1 and c_2. More precisely, let

$$c_1=oq_1^{-1}o_1^{-1}p_1, \quad \text{and} \quad c_2=oq_2o_2^{-1}p_2^{-1},$$

where $q_1=[v_1,z]$, $q_2=[z,v_2]$ are segments of q_0. To prove the proposition we need a few auxiliary lemmas.

Lemma 3.16. *Let $z\in V$, $o\in O(z)$, and let s be an H_i–component of the path o for a certain i. Then for $j=1,2$, there exist no H_i–components of o_j connected to s.*

Proof. First assume that s is connected to an H_i–component t of o_1 (see Fig. 3.2). This means that

$$dist_{X\cup\mathcal{H}}(s_-,t_-)\le 1.$$

Using (3.10), (3.11), and (3.7), we obtain

$$\begin{aligned} dist_{X\cup\mathcal{H}}(u,u_1) &\le dist_{X\cup\mathcal{H}}(u,s_-)+dist_{X\cup\mathcal{H}}(s_-,t_-)+dist_{X\cup\mathcal{H}}(t_-,u_1)\\ &\le (dist_{X\cup\mathcal{H}}(u,z)-1)+1+(l(o_1)-1)\\ &< K+K_0-1<2K. \end{aligned}$$



Figure 3.2:

By our choice of u_1 and v_1, this inequality implies $u_1 = p_-$, $v_1 = q_-$. Therefore, according to our choice of o_i, o_1 is a path in $\Gamma(G, X)$. Thus o_1 contains no H_i–components at all. A contradiction. The case $j = 2$ is completely analogous. □

For convenience, we give one more auxiliary definition. Given $z \in V$ and $o \in O(z)$, an H_i–component s of the path o is called an *ending component* if s contains z; for otherwise s is called a *non–ending* component. It can happen that o has no ending component, since the last edge of o can be labelled by a letter from X.

Lemma 3.17. *Let $z \in V$, $o \in O(z)$, and let s be a non–ending H_i–component of the path o for a certain i. Then for any $j = 1, 2$ the following holds. If s is not an isolated H_i–component of c_j, then there exists an H_i–component t of p_j that is connected to s.*

Proof. Again we consider the case $j = 1$ only. According to Lemma 3.16, there are no H_i–component of o_1 connected to s. Assume that there exists an H_i–component t of q_1 connected to s (see Fig. 3.3). Since s is a non–ending component, we have $dist_{X\cup\mathcal{H}}(s_-, z) \geq 2$. However, $dist_{X\cup\mathcal{H}}(s_-, t_-) \leq 1$. Hence

$$\begin{aligned} dist_{X\cup\mathcal{H}}(u, t_-) &\leq dist_{X\cup\mathcal{H}}(u, s_-) + dist_{X\cup\mathcal{H}}(s_-, t_-) \\ &< dist_{X\cup\mathcal{H}}(u, s_-) + dist_{X\cup\mathcal{H}}(s_-, z) \\ = dist_{X\cup\mathcal{H}}(u, z) \end{aligned}$$

This contradicts to the assumption that z is a closest vertex to u on q_0. □

From Lemma 3.17 we immediately obtain



Figure 3.3:

Corollary 3.18. *Let $z \in V$, $o \in O(z)$. Then for any $i = 1, \ldots, m$, every non–ending H_i–component of the path o is an isolated H_i–component of at least one of the cycles c_1, c_2.*

Proof. Indeed suppose that some non–ending H_i–component s of o is not isolated in both c_1 and c_2. By Lemma 3.17, this means that there are H_i–components t_1 and t_2 of p_1 and p_2 respectively that are connected to s. In particular, t_1 is connected to t_2. However this contradicts to the assumption that p is a locally minimal path without backtracking. □

Lemma 3.19. *Let $j = 1$ or $j = 2$. Suppose that for any $z \in V$ and any $o \in O(z)$, o has the ending component, which is not isolated in c_j; then for any $z \in V$ and any $o \in O(z)$, the ending component of o is connected to a component of p_j for corresponding i.*

Proof. For definiteness assume $j = 1$. We proceed by induction on $dist_{X\cup\mathcal{H}}(z, v_1)$. In case $dist_{X\cup\mathcal{H}}(z, v_1) = 0$, Lemma 3.19 is obvious. Indeed, assume that the ending component s of $o \in O(z)$ is not isolated in c_1. By Lemma 3.16, there are no components of o_1 connected to s. As o is a geodesic path, s is an isolated component of o. Therefore, the only possible case is that there exists a component of p_1 connected to s, since q_1 is trivial in this case.

Now assume that $dist_{X\cup\mathcal{H}}(z, v_1) \geq 1$. Arguing as above, we can easily show that it is sufficient to consider the case when s is connected to a component t of q_1. Let c be the edge of $\Gamma(G, X \cup \mathcal{H})$ labelled by a letter from $\widetilde{H}_i \setminus \{1\}$ for corresponding i such that $c_- = s_-$ and $c_+ = t_-$ (see Fig. 3.4). Since $dist_{X\cup\mathcal{H}}(u, t_-) = dist_{X\cup\mathcal{H}}(u, z)$, we have $t_- \in V$. Consider the new path $o' \in O(t_-)$ defined by $o' = [u, s_-]c$, where $[u, s_-]$ is the segment of the path o. Obviously the distance between t_- and v_1 is smaller that $dist_{X\cup\mathcal{H}}(v_1, z)$ and c is the ending component of o'. Thus, by the inductive assumption, there is an H_i–component r of p_1 connected to c. As c is connected to s, we get what we need. The lemma is proved. □



Figure 3.4:

Arguing as in the proof of Corollary 3.18, we immediately obtain

Corollary 3.20. *There exists a vertex $z \in V$ and a path $o \in O(z)$ such that either the last edge of o is labelled by a letter from X or the ending component of o is isolated in at least one of the cycles c_1, c_2.*

Now let us return to the proof of Proposition 3.15. By Corollaries 3.18 and 3.20, there exists a vertex z on q_0 that satisfies (3.11) and a (geodesic) path o connecting u and z such that for any i, any H_i–component of o is isolated in c_1 or in c_2. Applying Lemma 3.1, we obtain

$$dist_X(s_-, s_+) \le MLl(c_j) \tag{3.12}$$

for each H_i–component s of o. Let us estimate the length of c_j. Since p and q are (λ, c)–quasi–geodesics, we have the following bounds on the lengths of the paths p_j:

$$l(p_j) \le \lambda dist_{X\cup\mathcal{H}}(u_1, u) + c \le 2\lambda K + c. \tag{3.13}$$

Further,

$$\begin{aligned} l(q_j) \quad & \le l(q_0) \le \lambda dist_{X\cup\mathcal{H}}(v_1, v_2) + c \\ & \le \lambda(dist_{X\cup\mathcal{H}}(v_1, u_1) + dist_{X\cup\mathcal{H}}(u_1, u_2) + dist_{X\cup\mathcal{H}}(u_2, v_2)) + c \\ & \le 6\lambda K + c. \end{aligned} \tag{3.14}$$

Finally, by combining (3.7), (3.10), (3.11), (3.13), and (3.14), we obtain

$$l(c_j) \le l(p_j) + l(o) + l(q_j) + l(o_j) \le 8\lambda K + 2c + 2K. \tag{3.15}$$

Inequalities (3.12) and (3.15) imply

$$dist_X(u, z) \le dist_{X\cup\mathcal{H}}(u, z)ML \max_{j=1,2} l(c_j) \le KLM(8\lambda K + 2c + 2K). \tag{3.16}$$

It remains to assume ε to be equal to the right hand side of (3.16). □



Figure 3.5:

Proposition 3.15 allows one to show that if p and q are quasi–geodesic paths without backtracking in $\Gamma(G, X \cup \mathcal{H})$ with 'close' endpoints, then for any 'long' component s of p there exists a component t of q connected to s and, moreover, the X–distances between corresponding endpoints of s and t is 'small'.

Lemma 3.21. *For any $\lambda \geq 1$, $c \geq 0$, $k \geq 0$, there are $C = C(\lambda, c, k)$ and $D = D(\lambda, c, k)$ satisfying the following conditions. Let p and q be a pair of k–similar (λ, c)–quasi–geodesics in $\Gamma(G, X \cup \mathcal{H})$ such that p is a path without backtracking. Then for any $i = 1, \ldots, m$ and any H_i–component s of p satisfying the condition $dist_X(s_-, s_+) > C$, there exists an H_i–component t of q such that t is connected to s;*

Proof. We set
$$C = LM(1 + \lambda(2\varepsilon + 1) + c + 2\varepsilon), \tag{3.17}$$
where $\varepsilon = \varepsilon(\lambda, c, k)$ is provided by Proposition 3.15. Let s be an H_i–component of p such that $dist_X(s_-, s_+) > C$. By Proposition 3.15, there exist vertices w_1, w_2 on q such that $dist_X(s_-, w_1) \leq \varepsilon$ and $dist_X(s_+, w_2) \leq \varepsilon$. Since q is (λ, c)–quasi–geodesic, the length of the segment $q_0 = [w_1, w_2]$ of the path q satisfies the inequality

$$\begin{aligned} l(q_0) &\leq \lambda dist_{X\cup\mathcal{H}}(w_1, w_2) + c \\ &\leq \lambda(dist_{X\cup\mathcal{H}}(w_1, s_-) + dist_{X\cup\mathcal{H}}(s_-, s_+) + dist_{X\cup\mathcal{H}}(s_+, w_2)) + c \\ &\leq \lambda(2\varepsilon + 1) + c. \end{aligned} \tag{3.18}$$

We fix some paths r_1 and r_2 such that $(r_1)_- = s_-$, $(r_1)+ = w_1$, $(r_2)_- = s_+$, $(r_2)_+ = w_2$, and r_1, r_2 are geodesic paths in $\Gamma(G, X)$ (not in $\Gamma(G, X \cup \mathcal{H})$). By Proposition 3.15,
$$l(r_i) \leq \varepsilon, \quad i = 1, 2. \tag{3.19}$$

Notice that since the labels of r_1, r_2 consist of letters from X, r_1 and r_2 contain no H_i–components. Suppose that there exist no H_i–components of q_0 connected to s. Then s is an isolated H_i–component of the cycle

$$b = sr_2q_0^{-1}r_1^{-1}.$$

By Lemma 3.1 and inequalities (3.18), (3.19), we have

$$\begin{aligned} dist_X(s_-, s_+) \leq \quad & LMl(b) \leq LM(l(s) + l(q_0) + l(r_1) + l(r_2)) \\ & \leq LM(1 + \lambda(2\varepsilon + 1) + c + 2\varepsilon) = C, \end{aligned}$$

that contradicts to the assumption $dist_X(s_-, s_+) > C$. $\square$

The next lemma shows that if there exist two connected components s and t of two quasi–geodesics in $\Gamma(G, X \cup \mathcal{H})$ with 'close' endpoints, then the endpoints of s and t are 'close' even in case both components are 'short'.

Lemma 3.22. *For any $\lambda \geq 1, c \geq 0$ there exists a constant $D = D(\lambda, c, k)$ such that the following condition holds. Let p, q be a pair of k–similar (λ, c)–quasi–geodesics without backtracking in $\Gamma(G, X \cup \mathcal{H})$. Suppose that s and t are connected H_i–components of p and q respectively. Then*

$$\max\{dist_X(s_-, t_-),\ dist_X(s_+, t_+)\} \leq D.$$

Proof. Let $p_1 = [p_-, s_-]$ and $q_1 = [q_-, t_-]$ denote the corresponding segments of the paths p and q respectively, c' and c'' denote the connectors of length at most 1 such that $c'_- = s_-$, $c'_+ = t_-$, $c''_- = s_+$, and $c''_+ = t_+$ (see Fig. 3.6). Let $C(\lambda, c, k)$ be the constant provided by Lemma 3.21. We set

$$D = C(\lambda, c + \lambda + 1, k).$$

Let us estimate the X–length of c'.

Assume that the path p_1c' has a backtracking. Then there exists an H_i–component b of p_1 that is connected to c'. Therefore b is connected to s contradicting to the assumption that p is a path without backtracking. The same arguments show that no component of q_1 is connected to c'. Further, by Lemma 3.4 the path p_1c' is $(\lambda, c + \lambda + 1)$–quasi–geodesic. Thus we can apply Lemma 3.21 to q_1, p_1c', and the H_i–component c' of p_1c'. We obtain

$$dist_X(s_-, t_-) \leq C(\lambda, c + \lambda + 1, k)$$

The bound on $dist_X(s_+, t+)$ can be obtained in the same way. $\square$

Taking together Proposition 3.15 and Lemmas 3.21, 3.22, we obtain the following theorem, which is closely related to Farb's Bounded Coset Penetration property (see the appendix). For simplicity we change the notation and denote by $\varepsilon(\lambda, c, k)$ the maximal constant among $\varepsilon(\lambda, c, k)$ from Proposition 3.15, $C(\lambda, c, k)$ from Lemma 3.21, $D(\lambda, c, k)$ from Lemma 3.22.



Figure 3.6:

Theorem 3.23. *For any* $\lambda \geq 1$, $c \geq 0$, $k \geq 0$, *there exists a constant* $\varepsilon = \varepsilon(\lambda, c, k)$ *such that for any two* k*–similar* (λ, c)*–quasi–geodesics without backtracking* p *and* q *in* $\Gamma(G, X \cup \mathcal{H})$, *the following conditions hold.*

1) The sets of phase vertices of p *and* q *are contained in the closed* ε*–neighborhoods (with respect to the metric* $dist_X$*) of each other.*

2) Suppose that s *is an* H_i*–component of* p *such that* $dist_X(s_-, s_+) > \varepsilon$*; then there exists an* H_i*–component* t *of* q *which is connected to* s*.*

3) Suppose that s *and* t *are connected* H_i*–components of* p *and* q *respectively. Then*

$$\max\{dist_X(s_-, t_-),\ dist_X(s_+, t_+)\} \leq \varepsilon.$$

3.3 Geodesic triangles in Cayley graphs

Sometimes it is useful to think of δ–hyperbolic spaces as being fattened versions of trees (see [9], [27], and [44]). More precisely, given any three positive numbers a, b, c, we can consider the metric tree $T(a, b, c)$ that has three vertices of valence one, one vertex of valence three, and edges of length a, b, and c. For convenience we extend the definition of $T(a, b, c)$ in the obvious way to cover the cases where a, b and c are allowed to be zero.

Given any three point x, y, z in a metric space, the triangle inequality tells us that there exists (unique) non–negative numbers a, b, and c such that $dist(x, y) = a + b$, $dist(x, z) = a + c$, $dist(y, z) = b + c$. There is an isometry from $\{x, y, z\}$ to a subset of vertices of $T(a, b, c)$ (the vertices of valence one in the non–degenerate case); we label these vertices v_x, v_y, v_z in the obvious way (see Fig. 3.7).

For a geodesic triangle $\Delta = \Delta(x, y, z)$ with vertices x, y, z, we define $T_\Delta = T(a, b, c)$, where a, b, c are chosen as above. By o_Δ we denote the central vertex of T_Δ. The above map $\{x, y, z\} \to \{v_x, v_y, v_z\}$ extends uniquely to a map



Figure 3.7: The map χ_Δ. The points o_1, o_2, o_3 are mapped to o_Δ

$\chi_\Delta : \Delta \to T_\Delta$ whose restriction to each side of Δ is an isometry.

Definition 3.24. Let Δ be a geodesic triangle in a metric space Y. Consider the map $\chi_\Delta : \Delta \to T_\Delta$ defined above. We say that a point $u \in \Delta$ is *conjugate* to a point $v \in \Delta$ if $\chi_\Delta(u) = \chi_\Delta(v)$. The triangle Δ is said to be ξ*–thin* if $dist(u,v) \le \chi$ for any two conjugate points $u, v \in \Delta$.

The next lemma is well–known. It provides an equivalent definition of relative hyperbolicity.

Lemma 3.25. *A geodesic metric space Y is hyperbolic if and only if there exists $\xi \ge 0$ such that every geodesic triangle in Y is ξ–thin.*

The main result of this section is the relative analogue of the Rips condition for hyperbolic spaces. Namely we show that geodesic triangles in $\Gamma(G, X \cup \mathcal{H})$ are thin with respect to the metric $dist_X$ in the following sense.

Theorem 3.26. *There exists a constant $\nu > 0$ having the following property. Let $\Delta = pqr$ be a triangle whose sides p, q, r are geodesics in $\Gamma(G, X \cup \mathcal{H})$. Then for any vertex v on p, there exists a vertex u on the union $q \cup r$ such that*

$$dist_X(u,v) \le \nu.$$

Proof. The logical scheme of the proof is similar to that of Proposition 3.15. First we need an auxiliary construction. Let T_Δ be the tree related to Δ and $\chi_\Delta : \Delta \to T_\Delta$ be the corresponding map. Recall that $p_- = r_+, p_+ = q_-$. Let us consider vertices v_1, v_2 of p chosen as follows. Since $\Gamma(G, X \cup \mathcal{H})$ is a δ–hyperbolic space for some $\delta > 0$, according to Lemma 3.25, there exists a constant ξ such that Δ is ξ–thin. Increasing ξ if necessary we can assume $\xi \in \mathbb{N}$.

We chose vertices v_1 and v_2 on p in the following way. If $dist_{X\cup\mathcal{H}}(p_-, v) \le 6\xi$ (respectively $dist_{X\cup\mathcal{H}}(v, p_+) \le 6\xi$), we put $v_1 = p_-$ (respectively $v_2 = p_+$). If $dist_{X\cup\mathcal{H}}(p_-, v) > 6\xi$ (respectively $dist_{X\cup\mathcal{H}}(v, p_+) > 6\xi$), we take v_1 on the segment $[p_-, v]$ of the path p (respectively v_2 on the segment $[v, p_+]$ of the path p) such that

$$dist_{X\cup\mathcal{H}}(v, v_i) = 6\xi \tag{3.20}$$



Figure 3.8: The quadrilateral and hexagonal cases in the proof of Lemma 3.26.

for $i = 1$ (respectively $i = 2$).

Denote by u_1, u_2 the vertices on $q \cup r$ such that $\chi_\Delta(u_i) = \chi_\Delta(v_i)$, $i = 1, 2$. By Lemma 3.25, we have

$$dist_{X\cup\mathcal{H}}(v_i, u_i) \le \xi, \quad i = 1, 2. \tag{3.21}$$

Reversing the roles of p and r if necessarily, we can assume that $u_1 \in r$. Thus there are only two cases to consider (see Fig. 3.8).

Case 1 (Quadrilateral). The vertex u_2 lays on r. In this case we denote by p_1, p_2 (respectively by r_0) the segments of p (respectively the segment of r) such that $(p_1)_- = v_1$, $(p_1)_+ = v$, $(p_2)_- = v$, $(p_2)_+ = v_2$ (respectively $(r_0)_- = u_2$, $(r_0)_+ = u_1$). We also denote by o_1, o_2 the geodesic paths in $\Gamma(G, X \cup \mathcal{H})$ such that $(o_i)_- = v_i$, $(o_i)_+ = u_i$, $i = 1, 2$.

Case 2 (Hexagonal). No vertex on r is conjugate to v_2 (therefore, u_2 lays on q, see Fig. 3.8). We take a vertex w_1 on the segment $[r_-, u_1]$ of the path r such that $w_1 = r_-$ if $dist_{X\cup\mathcal{H}}(r_-, u_1) \le 12\xi$ and

$$dist_{X\cup\mathcal{H}}(w_1, u_1) = 12\xi \tag{3.22}$$

if $dist_{X\cup\mathcal{H}}(r_-, u_1) > 12\xi$. Furthermore, if $w_1 = r_-$, we set $w_2 = w_1$, and if $w_1 \ne r_-$, let w_2 be the vertex on $p \cup q$ such that $\chi_\Delta(w_2) = \chi_\Delta(w_1)$. Thus we have

$$dist_{X\cup\mathcal{H}}(w_1, w_2) \le \xi. \tag{3.23}$$

It is easy to see that $w_2 \in q$. Indeed, if $w_2 \in p$, then

$$dist_{X\cup\mathcal{H}}(v_1, w_2) = dist_{X\cup\mathcal{H}}(u_1, w_1) = 12\xi \ge dist_{X\cup\mathcal{H}}(v_1, v_2),$$

i.e., the point v_2 belongs to the segment $[v_1, w_2]$ of p. Since r contains conjugate points for both v_1 and w_2, there is a conjugate point for v_2 on r that contradicts to our assumption. Similar arguments show that w_2 belongs to the segment $[u_2, r_-]$ of the path q.

We denote by p_1, p_2 (respectively by r_0) the segments of p (respectively the segment of r) such that $(p_1)_- = v_1$, $(p_1)_+ = v$, $(p_2)_- = v$, $(p_2)_+ = v_2$ (respectively $(r_0)_- = w_1$, $(r_0)_+ = u_1$), and denote by q_0 the segment of q such that $(q_0)_- = u_2$, $(q_0)_+ = w_2$. Also let o_1, o_2, o_3 denote the geodesic paths in $\Gamma(G, X \cup \mathcal{H})$ such that $(o_i)_- = v_i$, $(o_i)_+ = u_i$, $i = 1, 2$, $(o_3)_- = w_1$, $(o_3)_+ = w_2$.

We deal with the second case in details, the first one is only slightly different. Let V denote the set of all vertices z on $r_0 \cup q_0$ such that

$$dist_{X\cup\mathcal{H}}(v, z) = \min_{z_0 \in r_0 \cup q_0} dist_{X\cup\mathcal{H}}(v, z_0), \tag{3.24}$$

where the minimum is taken among all vertices of $r_0 \cup q_0$. In particular, we have

$$dist_{X\cup\mathcal{H}}(v, z) \le \xi \tag{3.25}$$

since the conjugate point for z is a vertex on $r_0 \cup q_0$.

To every $z \in V$, we associate the set $O(z)$ of all a geodesic paths o such that $o_- = v, o_+ = z$. These paths cut the cycle $p_1 p_2 o_2 q_0 o_3^{-1} r_0 (o_1)^{-1}$ into two parts denoted by c_1 and c_2. More precisely, if $z \in r_0$, we set

$$c_1 = o r_1 (o_1)^{-1} p_1,$$

and

$$c_2 = o (r_2)^{-1} o_3 (q_0)^{-1} (o_2)^{-1} (p_2)^{-1},$$

where r_1 (respectively r_2) is the segment $[z, u_1]$ (respectively $[w_1, z]$) of the path r_0. If $z \in q_0$, we set

$$c_1 = o q_2 o_3^{-1} r_0 o_1^{-1} p_1$$

and

$$c_2 = p_2 o_2 q_1 o^{-1},$$

where q_1 (respectively q_2) is the segment $[u_2, z]$ (respectively $[z, w_2]$) of the path q_0. We emphasize that c_1, c_2 depend on the choice of $z \in V$ and $o \in O(z)$.

Lemma 3.27. *Let $z \in V$, $o \in O(z)$, and let s be an H_i–component of the path o for a certain i. Then for any $j = 1, 2, 3$ there exist no H_i–components of o_j connected to s.*

Proof. Without loss of generality we may assume that $z \in r_0$; the case $z \in q_0$ can be treated in the similar way and we leave it to the reader.

The proof of the lemma in the cases $j = 1$ and $j = 2$ almost coincide with the proof of the Lemma 3.16. Indeed assume that s is connected to an H_i–component t of o_j for $j = 1$ or $j = 2$. This means that

$$dist_{X \cup \mathcal{H}}(s_-, t_-) \leq 1.$$

Thus we have

$$\begin{aligned} dist_{X \cup \mathcal{H}}(v, v_j) &\leq dist_{X \cup \mathcal{H}}(v, s_-) + dist_{X \cup \mathcal{H}}(s_-, t_-) + dist_{X \cup \mathcal{H}}(t_-, v_j) \\ &\leq (\xi - 1) + 1 + (\xi - 1) < 2\xi. \end{aligned}$$

By our choice of v_j, this implies $v_j = p_-$ in case $j = 1$ and $v_j = p_+$ in case $j = 2$. Therefore, v_j coincides with u_j as

$$dist_{X \cup \mathcal{H}}(v_1, p_-) = dist_{X \cup \mathcal{H}}(u_1, p_-)$$

and

$$dist_{X \cup \mathcal{H}}(v_2, p_+) = dist_{X \cup \mathcal{H}}(u_2, p+).$$

Thus o_j is trivial. A contradiction.

Suppose that s is connected to an H_i–component t of o_3. In particular this means that o_3 is non–empty. By the choice of w_1 and w_2 this implies the equality $dist_{X \cup \mathcal{H}}(u_1, w_1) = 12\xi$. Note that

$$dist_{X \cup \mathcal{H}}(u_1, z) \leq dist_{X \cup \mathcal{H}}(u_1, v_1) + dist_{X \cup \mathcal{H}}(v_1, v) + dist_{X \cup \mathcal{H}}(v, z) \leq 8\xi.$$



Figure 3.9: Two cases in the proof of Lemma 3.27.

On one hand, we obtain

$$dist_{X\cup\mathcal{H}}(z,w_1) \geq dist_{X\cup\mathcal{H}}(u_1,w_1) - dist_{X\cup\mathcal{H}}(u_1,z) \geq 12\xi - 8\xi = 4\xi. \quad (3.26)$$

On the other hand, we have

$$\begin{aligned} dist_{X\cup\mathcal{H}}(z,w_1) &\leq dist_{X\cup\mathcal{H}}(z,s_+) + dist_{X\cup\mathcal{H}}(s_+,t_-) + dist_{X\cup\mathcal{H}}(t_-,w_1) \\ &\leq (\xi-1)+1+(\xi-1) < 2\xi \end{aligned}$$

that contradicts to (3.26). The lemma is proved. □

As in the previous section, given $z \in V$ and $o \in O(z)$, an H_i–component s of the path o is called an *ending component* if s contains z; for otherwise s is called a *non–ending* component. The proof of the next lemma is completely analogous to the proof of Lemma 3.17. We leave details to the reader.

Lemma 3.28. *Let $z \in V$, $o \in O(z)$, and s be a non–ending H_i–component of the path o for a certain i. For any $j = 1, 2$, if s is not an isolated H_i–component of c_j, then there exists an H_i–component t of p_j that is connected to s.*

Arguing as in the proof of Corollary 3.18, we immediately obtain the following.

Corollary 3.29. *Let $z \in V$, $o \in O(z)$. Then for any $i = 1, \ldots, m$, every non–ending H_i–component of the path o is an isolated H_i–component of at least one of the cycles c_1, c_2.*

The next lemma is the analogue of Lemma 3.19.

Lemma 3.30. *For any $j = 1, 2$, the following assertion is true. Suppose that for any $z \in V$ and any $o \in O(z)$, o has the ending component, which is not isolated in c_j; then for any $z \in V$ and any $o \in O(z)$, the ending component of o is connected to an H_i–component of p_j for corresponding i.*

Proof. For definiteness assume $j = 1$. For any $z \in V$, we define a non-negative integer number $\pi(z)$ as follows

$$\pi(z) = \begin{cases} dist_{X\cup\mathcal{H}}(u_1,z), \text{ if } z \in r_0, \\ \\ dist_{X\cup\mathcal{H}}(u_1,w_1) + dist_{X\cup\mathcal{H}}(w_2,z), \text{ if } z \in q_0. \end{cases}$$

The reader will have no difficulties in proving this lemma in the same way as Lemma 3.19. The only difference is that we have to proceed by induction on $\pi(z)$. □

Obviously Lemma 3.30 yields

Corollary 3.31. *There exists a vertex $z \in V$ and a path $o \in O(z)$ such that either the last edge of o is labelled by a letter from X or the ending component of o is isolated in at least one of the cycles c_1, c_2.*

Let us return to the proof of Theorem 3.26.

By Corollaries 3.29 and 3.31, there exists a vertex $z \in V$ and a path $o \in O(z)$ such that every component of o is isolated in at least one of the cycles c_1, c_2. Therefore, we have

$$dist_X(u, z) \leq l(o)ML \max\{l(c_1),\ l(c_2)\}. \tag{3.27}$$

Since

$$l(o) \leq \xi \tag{3.28}$$

by our choice of z, it remains to estimate the lengths of c_1 and c_2. Evidently we have

$$l(q_0) \leq l(o_3) + l(r_0) + l(o_1) + l(p_1) + l(p_2) + l(o_2) \leq 27\xi.$$

Therefore,

$$l(c_i) < l(o) + l(p_1) + l(p_2) + l(o_2) + l(q_0) + l(o_3) + l(r_0) + l(o_1) \leq 55\xi. \tag{3.29}$$

In view of (3.27), (3.28), and (3.29), to complete the proof it suffices to set $\nu = 55ML\xi^2$. □

Remark 3.32. For metric spaces, the Rips condition can be regarded as the definition of hyperbolicity. We note that the fulfilment of Theorem 3.26 for a given group and a collection of subgroups does not imply the relative hyperbolicity. Indeed for the pair $G \cong \mathbb{Z}$ and $H \cong 2\mathbb{Z}$ with the natural embedding $H \to G$, the statement of Theorem 3.20 obviously holds. However, G is not hyperbolic relative to H. Moreover, the corresponding relative Dehn function is not well–defined as follows from Proposition 2.36.

By drawing the diagonal, we obtain the following corollary of Theorem 3.20. It will be used in the next chapter to study the root problem for relatively hyperbolic groups.

Corollary 3.33. *Let $p_1p_2p_3p_4$ be a geodesic quadrangle in $\Gamma(G, X \cup \mathcal{H})$. Then for any vertex $u \in p_1$ there is a vertex $v \in p_2 \cup p_3 \cup p_4$ such that*

$$dist_X(u, v) \leq 2\nu.$$

3.4 Symmetric geodesics

The proofs of a number of theorems about ordinary hyperbolic groups (in particular, the solution of the conjugacy problem) are based on the following well–known property of hyperbolic metric spaces [19, 35]. Let p, q be two geodesics such that the distances between p_-, q_- and between p_+, q_+ are 'small', say less than or equal to 1, and u, v are two points on p and q respectively such that $dist(p_-, u) = dist(q_-, v)$. Then the distance between u and v is not greater than k, where k is independent of p, q, u, and v. The straightforward relative analogue of this property in the spirit of Proposition 3.15 and Theorem 3.26 can be stated as follows.

Conjecture 3.34. *For any* $k \geq 0$*, there exists a constant* $\varkappa = \varkappa(k)$ *such that the following condition holds. Let* p, q *be a pair of* k*–similar geodesics in* $\Gamma(G, X \cup \mathcal{H})$*. Then for any two vertices* $u \in p$ *and* $v \in q$ *such that* $dist(p_-, u) = dist(q_-, v)$*, we have*

$$dist_X(u, v) \leq \varkappa. \tag{3.30}$$

Unfortunately in general Conjecture 3.34 is false. Indeed consider the free group $G = \langle x, y\rangle$ and the subgroup $H = \langle x\rangle$, which is a free factor of G. Evidently G is hyperbolic relative to H. Let us also consider two geodesics, denoted by p and q, in the corresponding relative Cayley graph such that

$$\phi(p) \equiv yx^n, \quad \phi(q) \equiv x^n,$$

and

$$p_- = y^{-1}, \quad q_- = 1.$$

It is clear that p and q are 1–similar. Take the vertices $u \in p$ and $v \in q$ such that $dist_{X\cup\mathcal{H}}(p_-, u) = dist_{X\cup\mathcal{H}}(q_-, v) = 1$. Thus $u = 1$, $v = x^n$, and the length $u^{-1}v$ with respect to the generating set $\{x, y\}$ of G equals n. As n can be taken arbitrary large, this obviously violates (3.30).

However a certain refined version of the above mentioned conjecture can be proved for relatively hyperbolic groups. The results of this section will be used in the next chapter in order to study cyclic subgroups of relatively hyperbolic groups. We begin with definitions.

Definition 3.35. Let p, q be two paths in $\Gamma(G, X \cup \mathcal{H})$. We say that the pair (p, q) is *symmetric* if the labels of p and q coincide, i.e., $\phi(p) \equiv \phi(q)$. To each such a pair we associate the pair of elements $g_1 = (p_-)^{-1}q_-$, and $g_2 = (p_+)^{-1}q_+$ of G, called the *characteristic elements* of (p, q).

It is easy to see that two elements g_1, g_2 are conjugate by some element $t \in G$, i.e.,

$$g_2 = t^{-1}g_1t$$

if and only if there exists a symmetric pair of geodesics (p, q) in $\Gamma(G, X \cup \mathcal{H})$ such that (g_1, g_2) is the characteristic pair of (p, q) and $\overline{\phi(p)} = \overline{\phi(p)} = t$.

Definition 3.36. Let (p, q) be a pair of symmetric paths in $\Gamma(G, X \cup \mathcal{H})$. We say that two vertices $u \in p$, and $v \in q$ are *synchronous*, if $l([p_-, u]) = l([q_-, v])$, where $[p_-, u]$ and $[q_-, v]$ are segments of p and q respectively. Similarly, if a and b are H_i–components of p and q respectively for a certain i, we say that a and b are *synchronous components* when the vertices a_- and b_- are synchronous.

Now we are ready to state the main result of this section.

Theorem 3.37. *For any* $k \geq 0N$*, there exists a constant* $\varkappa = \varkappa(k)$ *with the following property. Let* (p, q) *be a symmetric pair of* k*–similar geodesics in* $\Gamma(G, X \cup \mathcal{H})$*,* u, v *a pair of synchronous vertices on* p *and* q *respectively. Then*

$$dist_X(u, v) \leq \varkappa.$$

The proof provided below can be slightly simplified by using references to some technical lemmas from [20]. However for convenience of the reader we give a complete proof in the spirit of our paper.

As usual, we divide the proof into a sequence of lemmas. The following is a particular case of a well–known property of hyperbolic spaces (see for example, [19]).

Lemma 3.38. *For any $k \geq 0$, there exists a constant $E = E(k)$ such that the following condition holds. Let p, q be a pair of k–similar geodesics in $\Gamma(G, X \cup \mathcal{H})$, and u, v synchronous vertices on p and q respectively. Then $dist_{X\cup\mathcal{H}}(u, v) \leq E$.*

We note that the constant E can be chosen effectively for a given k. In fact, one can take $E(k) = E'k$ for an appropriate constant E', which depends only on the hyperbolicity constant of $\Gamma(G, X \cup \mathcal{H})$, not on k.

Definition 3.39. We say that a symmetric pair of geodesics (p, q) in $\Gamma(G, X\cup\mathcal{H})$ is *minimal*, if for any other symmetric pair of geodesics (p', q') in $\Gamma(G, X \cup \mathcal{H})$ having the same characteristic elements, the inequality $l(p) \leq l(p')$ holds.

Lemma 3.40. *Let (p, q) be a minimal pair of symmetric geodesics in $\Gamma(G, X \cup \mathcal{H})$.*

1) Suppose that, for some i, two H_i–components a and b of p and q respectively are connected. Then a and b are synchronous.

2) Let u_1, v_1 and u_2, v_2 be two pairs of synchronous vertices of p and q respectively. Then $(u_1)^{-1}v_1 \neq (u_2)^{-1}v_2$.

Proof. 1) Suppose that a and b are not synchronous. Let

$$p = p_1 a p_2,$$

$$q = q_1 b q_2.$$

Assume for definiteness that

$$l(q_1) < l(p_1). \tag{3.31}$$

Since a and b are connected, there is an edge e in $\Gamma(G, X\cup\mathcal{H})$ such that $\phi(e) \in \widetilde{H}_i$ and

$$e_- = b_-, e_+ = a_+.$$

Also denote by r_1, r_2 some paths in $\Gamma(G, X \cup \mathcal{H})$ such that $(r_1)_- = p_-$, $(r_1)_+ = q_-$, $(r_2)_- = p_+$, $(r_2)_+ = q_+$ (see Fig. 3.10). Consider the cycles

$$c_1 = e^{-1} q_1^{-1} r_1^{-1} p_1 a$$

and

$$c_2 = e p_2 r_2 q_2^{-1} b^{-1}.$$



Figure 3.10: Gluing and cutting diagrams in the proof of Lemma 3.40.



Figure 3.11: The elements represented by the labels of the corresponding paths are written in brackets.

By Δ_1, Δ_2 we denote certain van Kampen diagrams over (3.1) whose boundaries have the same labels as c_1 and c_2 respectively. For simplicity, we will identify $\partial\Delta_j$ with c_j. Gluing Δ_1 and Δ_2 together along subpaths e^{-1} and e of their boundaries, we obtain a new diagram Δ over (3.1) with the boundary $r_1^{-1}pr_2q^{-1}$. Further by gluing together the subpaths p and q (which have the same labels), we transform Δ into an annular diagram Ξ (Fig. 3.10). Finally, we cut Ξ along the image of the path q_1ep_2 in Ξ and obtain a new diagram Σ over (3.1) with the boundary label

$$\phi(\Sigma) \equiv \phi(r_1)^{-1}U\phi(r_2)U^{-1},$$

where

$$U \equiv \phi(q_1ep_2).$$

Thus

$$g_2 = \overline{U}^{-1}g_1\overline{U} \tag{3.32}$$

in the group G, where $g_1 = \overline{\phi(r_1)}$, $g_2 = \overline{\phi(r_2)}$ are the characteristic elements of (p,q). The equality (3.32) leads to the symmetric pair (p',q') of geodesics in $\Gamma(G, X \cup \mathcal{H})$ such that (g_1,g_2) is the characteristic pair of (p',q') and $\phi(p') \equiv \phi(q') \equiv U$. According to (3.31), we have

$$l(p') = \|U\| = l(q_1) + 1 + l(p_2) < l(p_1) + 1 + l(p_2) = l(p)$$

that contradicts to the assumption that the pair (p,q) is minimal.

2) Let $(u_1)^{-1}v_1 = (u_2)^{-1}v_2 = w$ and $p = t_1t_2t_3$, where $t_1 = [p_-, u_1]$, $t_2 = [u_1, u_2]$, $t_3 = [u_2, p_+]$. Denote by f_1, f_3 the elements represented by labels of t_1, t_3 respectively (see Fig. 3.11). Then we have the following equalities in the group G: $g_2 = f_3^{-1}wf_3$, $w = f_1^{-1}g_1f_1$, where g_1, g_2 are the characteristic elements of (p,q). Thus g_1, g_2 are conjugate by the element f_1f_3 whose length is smaller than $l(p)$. We get a contradiction with the minimality of (p,q) again. $\square$

It is worth to note that the condition (3.33) in the following lemma is weaker than k–similarity of p and q. (This is important for our goals.)

Lemma 3.41. *Let (p,q) be a minimal pair of symmetric geodesics in $\Gamma(G, X \cup \mathcal{H})$ such that*

$$\max\{dist_{X\cup\mathcal{H}}(p_-, q_-),\ dist_{X\cup\mathcal{H}}(p_+, q_+)\} \le k \tag{3.33}$$

and let u and v be synchronous vertices on p and q respectively such that

$$\min\{dist_{X\cup\mathcal{H}}(p_-, u),\ dist_{X\cup\mathcal{H}}(u, p_+)\} \ge 2E,$$

where $E = E(k)$ is the constant provided by Lemma 3.38. Then

$$dist_X(u,v) \le 6MLE^2k.$$



Figure 3.12:

Proof. We repeat the trick used in the proof of Proposition 3.15. By the conditions of our lemma, there exist vertices $u_1, u_2 \in p$ and $v_1, v_2 \in q$ such that $u_1 \neq u_2$, $v_1 \neq v_2$ and

$$dist_{X\cup\mathcal{H}}(u_i, u) = dist_{X\cup\mathcal{H}}(v_i, v) = 2E. \tag{3.34}$$

For definiteness we assume that going along p (respectively along q) we first meet u_1 (respectively v_1) and then u_2 (respectively v_2).

Let o_i be a geodesic path in $\Gamma(G, X\cup\mathcal{H})$ such that $(o_i)_- = u_i$, $(o_i)_+ = v_i$ for $i = 1, 2$, and o be a geodesic path in $\Gamma(G, X\cup\mathcal{H})$ such that $o_- = u$, $o_+ = v$. We consider the cycles

$$c_1 = [v_1, v]o^{-1}[u_1, u]^{-1}o_1$$

and

$$c_2 = [v, v_2]o_2^{-1}[u, u_2]^{-1}o,$$

where $[u_1, u]$ and $[u, u_2]$ (respectively $[v_1, v]$ and $[v, v_2]$) are the segments of p (respectively q).

Note that every component of o is an isolated component in at least one of the cycles c_1, c_2. Indeed, the same arguments as in the proof of Lemma 3.16 together with the equalities (3.34) show that no component of o_i is connected to a component of o for $i = 1, 2$. Suppose that a component s of the path o is connected to a component r_1 in c_1 and r_2 in c_2. Since p and q are geodesics, r_1 and r_2 can not belong simultaneously to p or q. Hence we can assume that $r_1 \in p$ and $r_2 \in q$ (see Fig. 3.12). By Lemma 3.40, r_1 and r_2 are synchronous components. However this can not happen if $r_1 \in c_1$ and $r_2 \in c_2$.

Thus every component of o is an isolated component in at least one of the cycles c_1, c_2 and has X–length at most $ML\max\limits_{i=1,2} l(c_i)$ by Lemma 3.1. By Lemma 3.38, $l(o_i) \leq E$ for $i = 1, 2$ and $l(o) \leq E$. Therefore, $l(c_i) \leq 6E$ for $i = 1, 2$. This implies

$$dist_X(u, v) \leq l(o)ML\max_{i=1,2} l(c_i) \leq 6MLE^2.$$

$\square$

Corollary 3.42. *For any $k \geq 0$ there exists a constant $\rho = \rho(k)$ such that for any two conjugate elements f, g of G of relative lengths*

$$\max\{|f|_{X\cup\mathcal{H}},\ |g|_{X\cup\mathcal{H}}\} \leq k,$$

there exists an element $t \in G$ such that $f^t = g$ and

$$|t|_{X\cup\mathcal{H}} \leq \rho.$$

Proof. Let

$$\rho = (card\, X)^{6MLE^2k} + 1 + 4E,$$

where $E = E(k)$ is the constant from Lemma 3.38. Let (p, q) be the minimal pair of symmetric geodesics in $\Gamma(G, X\cup\mathcal{H})$ with the characteristic elements f, g. Set $t = \overline{\phi(p)}$. If $l(p) > 4E$, let p_0 be the segment of p such that

$$dist_{X\cup\mathcal{H}}((p_0)_-, p_-) = dist_{X\cup\mathcal{H}}((p_0)_+, p_+) = 2E.$$

By the second statement of Lemma 3.40 and Lemma 3.41, the length of p_0 does not exceed the number of elements of G having length at most $6MLE^2k$ with respect to the generating set X. Thus we have

$$l(p_0) \leq (card\, X)^{6MLE^2k} + 1$$

and

$$|t|_{X\cup\mathcal{H}} = l(p) \leq l(p_0) + 4E \leq \rho.$$

□

Lemma 3.43. *For any $k \geq 0$, $\lambda \geq 1$, $c \geq 0$, there exists a constant $\eta = \eta(\lambda, c, k)$ such that the following condition holds. Let (p, q) be an arbitrary symmetric pair of k–similar (λ, c)–quasi-geodesics without backtracking in $\Gamma(G, X\cup\mathcal{H})$ such that no synchronous components of p and q are connected. Then for any $i = 1, \ldots, m$ and any H_i–component e of p, we have*

$$dist_X(e_-, e_+) \leq \eta l([p_-, e_+]), \tag{3.35}$$

where $[p_-, e_+]$ is the segment of p.

Proof. Let $\varepsilon_0 = \varepsilon(\lambda, c, k)$ be the constant given by Theorem 3.23. Let also $\varepsilon_1 = \varepsilon(\lambda, c, \max\{k, \varepsilon_0\})$. Set

$$\eta = \max\{2\varepsilon_0, \varepsilon_1, 1\}.$$

We proceed by induction on $n = l([p_-, e_+])$. The proof in the case $n = 1$ is given below together with the proof in the general case.

First suppose that no component of q is connected to e. Then the inequality (3.35) follows from Theorem 3.23. Further, assume that there exists an H_i–component e' of q connected to e. Let m be the length of the segment $[q_-, e'_+]$



Figure 3.13: The cases $m > n$ and $m < n$ in the proof of Lemma 3.43.

of q. By our assumption, $m \neq n$. Thus there are two possibilities (see Fig. 3.13).

Case 1. $m < n$ (this case is impossible if $n = 1$). Note that

$$\max\{dist_X(e_-, e'_-),\ dist_X(e_+, e'_+)\} \leq \varepsilon_0 \tag{3.36}$$

by Theorem 3.23. We denote by f the H_i–component of p which is synchronous to e'. Since p and q are symmetric, we have

$$dist_X(e'_-, e'_+) = dist_X(f_-, f_+) \leq \eta m \tag{3.37}$$

by the the inductive assumption. Taking (3.36) and (3.37) together, we obtain

$$dist_X(e_-, e_+) \leq 2\varepsilon_0 + \eta m \leq \eta(m+1) \leq \eta n$$

Case 2. $m > n$. Denote by g the H_i–component of q which is synchronous to e. If no H_i–component of the segment $[p_-, e_-]$ of p is connected to g (in particular, this is so if $n = 1$ since the segment $[p_-, e_-]$ is trivial in this case), then we obtain

$$dist_X(e-, e_+) = dist_X(g_-, g_+) \leq \varepsilon_1$$

by applying Theorem 3.23 for the segments $[p_-, e_-]$ and $[q_-, e'_-]$ of p and q. The case when g is connected to a component of the segment $[p_-, e_-]$ can be reduced to *Case 1* by reversing the roles of p and q. $\square$



Figure 3.14:

Lemma 3.44. *For any $k \in \mathbb{N}$, there exists a constant $\varkappa_0 = \varkappa_0(k)$ such that the following condition holds. Let (p, q) be a symmetric pair of k–similar geodesics in $\Gamma(G, X \cup \mathcal{H})$ such that no synchronous components of p and q are connected. Then for every pair of synchronous vertices $u \in p$, $v \in q$, we have*

$$dist_X(u, v) \leq \varkappa_0.$$

Proof. Without loss of generality, we may assume that $q_- = 1$. Since the labels of the segments $[p_-, u]$ and $[1, v]$ of p and q respectively coincide, the element p_- is conjugate to the element $w = v^{-1}u$ in G.

Note that by Lemma 3.38, the relative length of w satisfies $|w|_{X\cup\mathcal{H}} \leq E$. We consider the shortest (with respect to the relative metric) element $t \in G$ such that $p_- = t^{-1}wt$. By Lemma 3.42, the we have

$$|t|_{X\cup\mathcal{H}} \leq \rho,$$

where $\rho = \rho(\max\{E, k\})$. Let (r, s) be the symmetric pair of geodesics in $\Gamma(G, X \cup \mathcal{H})$ such that $r_- = u$, $s_- = v$, and the labels $\phi(r) \equiv \phi(s)$ represent t in G. Thus the label of any path from s_+ to r_+ in $\Gamma(G, X \cup \mathcal{H})$ represents the element p_- in G.

Suppose that there exist components of r connected to some components of p. We consider the last such a component a of r. Thus $r = r_1 a r_2$, there exists a component b of p connected to a, and no component of r_2 is connected to a component of p. To be definite we assume that b belongs to $[u, p_+]$. (The case when b belongs to $[p_-, u]$ can be treated in the same way and is left to the reader.) Let c be a path in $\Gamma(G, X \cup \mathcal{H})$ of length at most 1 such that $c_- = b_-$, $c_+ = a_+$ (see Fig. 3.14). Notice that

$$l([u, b_-]) = l([u, b_+]) - 1 \leq dist_{X\cup\mathcal{H}}(u, a_-) + dist_{X\cup\mathcal{H}}(a_-, b_+) - 1 = l(r_1)$$

as p and r are geodesic. Hence

$$l([u,b_-]cr_2) \leq l(r_1) + 1 + l(r_2) = l(r).$$

Obviously the path $[p_-, b_-]cr_2$ is $(1,\rho)$–quasi–geodesic without backtracking as follows from the choice of a. Thus replacing r with $r' = [u, b_-]cr_2$ and s with the geodesic symmetric to r', we may assume that $[p_-, u]r$ has no backtracking.

Let $\varepsilon = \varepsilon(1, \rho, k)$ and $\eta = \eta(1, \rho, \max\{\varepsilon, k\})$ be constants provided be Theorem 3.23 and Lemma 3.43 respectively. We consider two cases.

Case 1. Suppose that there are no synchronous connected components of r and s. Applying Lemma 3.43 for the symmetric $(1,\rho)$–quasi–geodesic paths $r^{-1}[p_-, u]^{-1}$ and $s^{-1}[q_-, v]^{-1}$, we obtain that the X–length of any H_i–component e of r satisfies the inequality

$$dist_X(e_-, e_+) \leq \eta l(r) \leq \eta\rho.$$

Hence,

$$dist_X(u, r_+) \leq l(r)\eta\rho \leq \eta\rho^2.$$

Finally we have

$$dist_X(u,v) \leq dist_X(u, r_+) + dist_X(r_+, s_+) + dist_X(s_+, v) \leq 2\eta\rho^2 + k.$$

Case 2. Now assume that there is at least one pair of synchronous connected components in r and s. Let i, j be the connected synchronous components of r and s respectively such that there are no connected synchronous components of the segments $[u, i_-]$ and $[v, j_-]$ of r and s. By Theorem 3.23, we have $dist_X(i_-, j_-) \leq \varepsilon$. Therefore, the paths $[p_-, u][u, i_-]$ and $[1, v][v, j_-]$ form a symmetric $\max\{\varepsilon, k\}$–similar pair. Arguing as in the Case 1, we obtain

$$dist_X(u,v) \leq 2\eta\rho^2 + \varepsilon.$$

In both cases it suffices to set $\varkappa_0 = 2\eta\rho^2 + \max\{\varepsilon, k\}$. □

Now we are ready to prove the main result of this section.

Proof of Theorem 3.37. Let $a_1, \ldots, a_l$ be the set of all components of p that are connected to the corresponding synchronous components of q. We denote by b_j the component of q connected to a_j. Changing if necessary the order of enumeration of $a_1, \ldots, a_l$, we may assume that

$$p = p_1 a_1 \ldots p_l a_l p_{l+1},$$

$$q = q_1 b_1 \ldots q_l b_l q_{l+1},$$

where for any $j = 1, \ldots, l+1$, (p_j, q_j) is a symmetric pair of ε–similar geodesics for $\varepsilon = \varepsilon(1, 0, k)$ given by Theorem 3.23 (see Fig ...). Notice that there exist no connected synchronous components of p_j and q_j for any $j = 1, \ldots, l+1$. As u and v are synchronous vertices of p_j and q_j for a certain j, it suffices to set $\varkappa = \varkappa_0(\max\{\varepsilon, k\})$, where $\varkappa_0(\max\{\varepsilon, k\})$ is provided by Lemma 3.44. □



Figure 3.15: The decomposition of p and q in the proof of Theorem 3.37.

Chapter 4

Algebraic properties

4.1 Elements of finite order

Recall that the number of conjugacy classes of elements of finite order in any hyperbolic group is finite. A generalization of this result to the class of hyperbolic products of groups can be found in [72]. In this section we extend these results by proving the following.

Definition 4.1. Let G be a group hyperbolic with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$. An element $g\in G$ is called *parabolic* if it is conjugate to an element of one of the subgroups H_λ. Otherwise g is said to be *hyperbolic.*

Theorem 4.2. *Suppose G is a group hyperbolic with respect to a collection of subgroups $\{H_\lambda\}_{\lambda\in\Lambda}$. Then the number of conjugacy classes of hyperbolic elements of finite order in G is finite.*

It is well known that if G acts on a tree without inversions and with a compact quotient, then every element of finite order has a fixed point (see [77] or [33]). In other terms, if G is a fundamental group of a a graph of groups, then each element of finite order in G is conjugate to an element of one of the vertex groups. In combination with Splitting Theorem from Section 2.4, this shows that it suffices to prove the theorem in case G is finitely generated (and therefore the collection of subgroups is finite by Corollary 2.48). In the rest of this section we assume that G is generated by a finite set X and is hyperbolic relative to subgroups $\{H_1,\ldots,H_m\}$.

Recall, that a path in a metric space is said to be k–local geodesic if any its subpath of length at most k is geodesic. The following lemma is well known (see, for example, [19, CH. III.H, Theorem 1.13]).

Lemma 4.3. *Let r be a k–local geodesic in a δ–hyperbolic metric space for some $k>8\delta$. Then r is (λ,c)–quasi–geodesic for $\lambda=\frac{k+4\delta}{k-4\delta}\leq 3$ and $c=2\delta$.*



Figure 4.1:

Corollary 4.4. *Let g be an element of finite order in G. Then G is conjugate to an element of relative length at most $8\delta + 1$, where δ is the hyperbolicity constant for $\Gamma(G, X \cup \mathcal{H})$.*

Proof. We assume that g is a shortest element in the conjugacy class g^G and $|g|_{X\cup\mathcal{H}} > 8\delta + 1$. Let us take a shortest word $U \in (X \cup \mathcal{H})^*$ representing g and consider the path p_n such that $(p_n)_- = 1$, $\phi(p_n) \equiv U^n$, $n \in \mathbb{N}$. Since g is a shortest element in g^G, p_n is $(8\delta + 1)$–local geodesic in $\Gamma(G, X \cup \mathcal{H})$. Therefore, by Lemma 4.3, we have

$$|g^n|_{X\cup\mathcal{H}} = dist_{X\cup\mathcal{H}}(1, (p_n)_+) \geq \frac{1}{3}l(p_n) - 2\delta \geq \frac{1}{3}|n| - 2\delta.$$

Note that $1/3|n| - 2\delta \neq 0$ for any n big enough. Hence the order of g is infinite. □

In contrast to the case of ordinary hyperbolic groups, the above corollary does note imply the desired result since balls in $\Gamma(G, X \cup \mathcal{H})$ are, in general, not finite. The next lemma is the crucial ingredient of the proof of Theorem 4.2.

Lemma 4.5. *Suppose that a group G is generated by a finite set X and is hyperbolic relative to subgroups $\{H_1, \ldots, H_m\}$. Then there exists a constant B such that the following condition holds. Let f be a hyperbolic element of finite order in G such that f has smallest relative length among all elements of the conjugacy class f^G. Then*

$$|f|_X \leq B|f|^2_{X\cup\mathcal{H}}.$$

Proof. Let U be a shortest word in $(X \cup \mathcal{H})^*$ representing f in G. Let n be the order of f, l the relative length of f. We consider an arbitrary cycle p in $\Gamma(G, X \cup \mathcal{H})$ with the label U^n and take a subpath p_0 of p obtained as follows. If p is a cycle without backtracking, we set $p_0 = p$. Further suppose that there exists two connected components s_1 and s_2 of p. Let

$$p = as_1bs_2c.$$

Passing to another pair of connected components if necessary, we may assume that b is a path without backtracking and no component of b is connected to the component s_1, s_2. In this case we set $p_0 = b$.

Since $\phi(p_0)$ is a (cyclic) subword of U^n, we have

$$\phi(p_0) \equiv U_0^k V,$$

where U_0 is a cyclic shift of U and V is a cyclic subword of U of length less than l. It is easy to check that $k > 0$. Indeed if $k = 0$, then f is not the shortest element in the conjugacy class f^G.

Let e be a path in $\Gamma(G, X \cup \mathcal{H})$ of length at most 1 such that $e_- = (p_0)_-$ and $e_+ = (p_0)_+$ (see Fig. 4.1). We consider the cycle $c = p_0 e^{-1}$. According to our choice of p_0, c is a cycle without backtracking. We have

$$l(c) = l(p_0) + 1 \leq kl + \|V\| + 1 \leq (k+1)l.$$

Let W be an H_j–syllable in U. Then W is a label of at least k components of c. Since all components of c are isolated, applying Lemma 3.1 we obtain

$$k|\overline{W}|_X \leq MLl(c) \leq ML(k+1)l.$$

Hence,

$$|\overline{W}|_X \leq ML\frac{k+1}{k}l \leq 2MLl. \tag{4.1}$$

Finally, since the inequality (4.1) holds for any syllable W of U, we have $|\overline{U}|_X \leq 2MLl^2$. □

Proof of Theorem 4.2. By combining Lemmas 4.4 and 4.5, we obtain that each hyperbolic element of finite order in G is conjugate to an element of X–length at most $B(8\delta+1)^2$. Since G is locally finite with respect to the metric $dist_X$, this proves the theorem. □

As a corollary, we have

Corollary 4.6. *The set of orders of hyperbolic elements in G is finite.*

Corollary 4.7. *If G is residually finite and all subgroups H_λ are torsion free, then G is virtually torsion free, that is G contains a torsion free subgroup of finite index.*

Proof. Let $g_1, g_2, \ldots, g_k$ be elements of G such that each hyperbolic element of finite order is conjugate to one of them. Then there exists a normal subgroup N of finite index in G such that $g_i \notin N$ for all $i = 1, \ldots, k$. Thus N contains no hyperbolic elements of finite order. Hence N is torsion free. □

Note that the requirement of residual finiteness is essential.

Example 4.8. Let H be a finitely generated torsion free simple group. Let w be an arbitrary nontrivial element of H. Consider the group G given by the relative presentation

$$G = \langle x, H \mid x^2 = 1,\ x = [x, w][x, w^2] \ldots [x, w^n] \rangle.$$

It is easy to check that G is a quotient of the free product $\langle x \mid x^2 = 1\rangle * H$ by the relation $x = [x, w][x, w^2] \ldots [x, w^n]$ satisfying $C'(\lambda)$ hypothesis (as a relation over a free product) with $\lambda \to 0$ as $n \to \infty$. Therefore, by the Greendlinger Lemma [56], the relative Dehn function of G with respect to H is linear for all n big enough and thus G is hyperbolic relative to H. However, any subgroup N of finite index in G contains H (otherwise $N \cap H = \{1\}$ and hence G/N is infinite). Now the relation $x = [x, w][x, w^2] \ldots [x, w^n]$ implies that $x \in G$. Thus $N = G$, i.e., G contains no proper subgroups of finite index. In particular, G is not virtually torsion free.

4.2 Relatively quasi–convex subgroups

Our discussion in this section is stimulated by some ideas of Gromov [45] which were elaborated by Gersten and Short [43], Alonso and Bridson [3], and others (see [35] and references therein). Our main goal here is to introduce the (geometric) notion of a quasi–convex subgroup of a relatively hyperbolic group and to obtain some analogues of well–known theorems about quasi–convex subgroups in hyperbolic groups. In the next section, adopting an idea of Gersten and Short [43] to the relative case, we apply our results to the study of translation numbers. For the dynamical notion of quasi–convexity for convergence groups we refer to [12].

Definition 4.9. Let G be a group generated by a finite set X, $\{H_1, \ldots, H_m\}$ a collection of subgroups of G. A subgroup R of G is called relatively quasi–convex with respect to $\{H_1, \ldots, H_m\}$ (or simply *relatively quasi–convex* when the collection $\{H_1, \ldots, H_m\}$ is fixed) if there exists a constant $\sigma > 0$ such that the following condition holds. Let f, g be two elements of R, and p an arbitrary geodesic path from f to g in $\Gamma(G, X \cup \mathcal{H})$. Then for any vertex $v \in p$, there exists a vertex $w \in R$ such that

$$dist_X(u, w) \le \sigma.$$

Note that, without loss of generality, we may assume one of the elements f, g to be equal to the identity since both the metrics $dist_X$ and $dist_{X\cup\mathcal{H}}$ are invariant under the left action of G on itself.

It is easy to see that, in general, this definition depends on X. However in case of relatively hyperbolic groups we have

Proposition 4.10. *Let G be a group hyperbolic with respect to a collection of subgroups $\{H_1, \ldots, H_m\}$ and R a subgroup of G. Suppose that X_1, X_2 are two* *finite generating sets of G. Then Q is relatively quasi–convex with respect to X_1 if and only if it is relatively quasi–convex with respect to X_2.*

Proof. Let $\Gamma_1 = \Gamma(G, X_1 \cup \mathcal{H})$ and $\Gamma_2 = \Gamma(G, X_2 \cup \mathcal{H})$. For every $x \in X_1$, we fix a word W_x over X_2 representing x in G. To each path p in Γ_1, we assign a path in Γ_2 which starts and ends at the same elements as p and has label obtained from $\phi(p)$ by replacing x with W_x for every $x \in X_1$.

Suppose that R is quasi–convex with respect to X_2. Let r be an element of R, p a geodesic path in Γ_1 such that $p_- = 1$ and $p_+ = r$. We also take a vertex $u \in p$. Denote by q the path in Γ_2 corresponding to p, and by v the vertex corresponding to v (thus $u = v$ being considered as elements of G). By Proposition 2.8, q is $(\lambda, 0)$–quasi–geodesic for some constant λ which is independent of p. Moreover, since p is a path without backtracking, then obviously so is q.

By Theorem 3.23, q lies in the closed $\varepsilon = \varepsilon(\lambda, 0, 0)$–neighborhood (with respect to the metric $dist_{X_2}$) of the geodesic path s in Γ_2 with $s_- = 1$, $s_+ = r$. As R is quasi–convex with respect to X_2, s belongs to the closed σ–neighborhood (with respect to $dist_{X_2}$) of R, where σ is the quasi–convexity constant. Thus

$$dist_{X_2}(v, R) \leq \varepsilon + \sigma.$$

Applying Proposition 2.8 again, we obtain the upper bound on $dist_{X_1}(u, R)$, which is independent of u and p. Thus R is quasi–convex with respect to X_1. ☐

Definition 4.11. Let G be as in the Definition 4.9. A relatively quasi–convex subgroup R of G is called *strongly relatively quasi–convex* if the intersection $R \cap H_i^g$ is finite for any $g \in G$, $i = 1, \ldots, m$.

Definition 4.12. Recall that a map $\iota : (M_1, dist_1) \to (M_2, dist_2)$ between two metric spaces M_1 and M_2 with metrics $dist_1$ and $dist_2$ is called a *quasi–isometric embedding* if there exist $c_1, c_2 > 0$ such that for every two points $x, y \in M_1$ we have

$$\frac{1}{c_1} dist_1(x, y) - c_2 \leq dist_2(\iota(x), \iota(y)) \leq c_1 dist_1(x, y) + c_2.$$

Theorem 4.13. *Suppose that the group G is hyperbolic relative to the collection of subgroups $\{H_1, \ldots, H_m\}$. Let R be a subgroup of G. Then the following conditions are equivalent.*

1. *R is strongly relatively quasi–convex.*
2. *R is generated by a finite set Y and the natural map $(R, dist_Y) \to (G, dist_{X \cup \mathcal{H}})$ is a quasi–isometric embedding.*

Proof. For every $x, y \in G$ and every $i = 1, \ldots, m$ we consider the set

$$Z_{x,y,i} = \{xhy \mid h \in H_i\} \cap R.$$



Figure 4.2:

Also set

$$Z_0 = \{r \in R \mid |r|_X \le 2\sigma + 1\},$$

where σ is the quasi–convexity constant for R.

To prove the theorem we need two auxiliary lemmas.

Lemma 4.14. *Let*

$$B_\sigma^X = \{g \in G \mid |g|_X \le \sigma\}. \tag{4.2}$$

Then the subgroup R is generated by the set

$$Z = Z_0 \bigcup \left(\bigcup_{x,y \in B_\sigma^X} \bigcup_{i=1}^{m} Z_{x,y,i} \right).$$

Proof. Let r be an arbitrary element of R. We consider a geodesic p in $\Gamma(G, X \cup \mathcal{H})$ such that $p_- = 1$, $p_+ = r$. Let $g_0 = 1, g_1, \dots, g_n = r$ be the consecutive vertices of p (see Fig. . By the definition of a relatively quasi–convex subgroup, for any $i = 1, \dots, n-1$, there exists an element $r_i \in R$ such that

$$dist_X(r_i, g_i) \le \sigma. \tag{4.3}$$

We also set $r_0 = 1$ and $r_n = r$. Denote by x_i the element $r_i^{-1} g_i$ and by e_{i+1} the edge of p going from g_i to g_{i+1}.

Obviously we have

$$r_{i+1} = r_i x_i \overline{\phi(e_{i+1})} x_{i+1}^{-1}.$$

By (4.3), the X–length of x_i satisfies $|x_i|_X \le \sigma$. Therefore the element

$$s_i = x_i \overline{\phi(e_{i+1})} x_{i+1}^{-1}$$

either belongs to Z_0 in case $\phi(e_{i+1}) \in X$, or belongs to $Z_{x_i, x_{i+1}, j}$ if $\phi(e_{i+1}) \in \widetilde{H}_j$ for some $j = 1, \dots, m$. Thus in both cases we have $s_i \in Z$.

It is clear that

$$r_n = r_{n-1} s_{n-1} = \ldots = r_0 s_0 \ldots s_{n-1} = s_0 \ldots s_{n-1} \tag{4.4}$$

as $r_0 = 1$. Therefore $r_n \in \langle Z \rangle$. Since r was an arbitrary element of R, the lemma is proved. $\square$

Lemma 4.15. *For any $x, y \in G$ and any $i = 1, \ldots, m$ the set $Z_{x,y,i}$ is finite.*

Proof. Assume that $Z_{x,y,i} = \{z_0, z_1, \ldots\}$ is infinite for some $x, y \in G$, $i \in \{1, \ldots, m\}$. Suppose that for any $j = 0, 1, \ldots$, $z_j = xh_jy$, where $h_j \in H_i$. The elements $t_j = z_0^{-1}z_j$ are different for different $j \geq 0$. However we have

$$t_j = y^{-1}h_0x^{-1}xh_jy = y^{-1}h_0h_jy \in H_i^y$$

for every j. Recall that $t_j \in R$ for any j by the definition of $Z_{x,y,i}$. Therefore, the intersection $R \cap H_i^y$ is infinite contrary to our assumption. $\square$

Now we are able to show that the first condition in Theorem 4.13 implies the second one. Using Lemma 4.15, one can easily see that the set Z is finite since the ball B_σ^X is finite. Further by Lemma 4.14, R is generated by Z. Therefore R is finitely generated. Moreover, it follows from the proof of Lemma 4.14 (see (4.4)) that for any element $r \in R$ of relative length $|r|_{X \cup \mathcal{H}} = n$, we have $r = s_0 \ldots s_{n-1}$ for certain $s_0, \ldots, s_{n-1} \in Z$. This means that

$$|r|_Z \leq |r|_{X \cup \mathcal{H}},$$

i.e., the map $(R, dist_Z) \to (G, dist_{X \cup \mathcal{H}})$ is a quasi–isometric embedding.

To prove the converse implication, we assume that R is generated by a finite set $Y = Y^{-1}$ and the natural map $(R, dist_Y) \to (G, dist_{X \cup \mathcal{H}})$ is a quasi–isometric embedding. We denote by c_1, c_2 the corresponding quasi–isometry constants.

Note that if $r \in R \cap H_i^g$ for some $i \in \{1, \ldots, m\}$ and $g \in G$, then

$$|r|_Y \leq c_1|r|_{X \cup \mathcal{H}} + c_2 \leq c_1(2|g|_{X \cup \mathcal{H}} + 1) + c_2.$$

Since Y is finite, we obtain $card\,(R \cap H_i^g) < \infty$ for $i = 1, \ldots, m$ and any $g \in G$.

It remains to show that R is relatively quasi–convex. For any element $y \in Y$, we fix a word W_y over the alphabet $X \cup \mathcal{H}$ representing y in G. Set

$$\mu = \max_{y \in Y} \|W_y\|. \tag{4.5}$$

Given an element $r \in R$, we consider the shortest word $V = y_1 \ldots y_n$ over Y representing r. Let

$$U = W_{y_1} \ldots W_{y_n}$$

be the word over X obtained from V by replacing each y_i with the corresponding W_{y_i}.

Let U_0 be a subword of U. Then

$$U_0 \equiv AW_{y_j} \ldots W_{y_{j+k}}B,$$

where $\|A\| < \mu$ and $\|B\| < \mu$. Since any subword of V is geodesic with respect to the metric $dist_Y$ on R, we have

$$\begin{aligned}\|U_0\| < \ & 2\mu + \mu(k+1) = \\ & 2\mu + \mu|y_j \dots y_{j+k}|_Y \leq \\ & 2\mu + \mu\left(c_1|y_j \dots y_{j+k}|_{X\cup\mathcal{H}} + c_2\right) \leq \\ & 2\mu + \mu\left(c_1\left(|\overline{U_0}|_{X\cup\mathcal{H}} + 2\mu\right) + c_2\right).\end{aligned}$$

Thus the path p in $\Gamma(G, X\cup\mathcal{H})$ with $p_- = 1$ labelled U is $(\mu c_1, 2\mu + 2\mu^2 c_1 + \mu c_2)$–quasi–geodesic. Let $\varepsilon = \varepsilon(\mu c_1, 2\mu + 2\mu^2 c_1 + \mu c_2, 0)$ be the constant provided by Proposition 3.15. Then for any geodesic path q in $\Gamma(G, X\cup\mathcal{H})$ such that $q_- = 1$ and $q_+ = r$, and any vertex $v \in q$, there exists a vertex $u \in p$ such that

$$dist_X(u, v) \leq \varepsilon.$$

It is clear that any vertex of p belongs to the closed μ–neighborhood of R with respect to the metric $dist_X$. Hence,

$$dist_X(v, R) \leq \varepsilon + \mu.$$

Since the right hand side of the above inequality is independent of r, R is relatively quasi–convex. □

As is well known, any quasi–convex subgroup of a hyperbolic group is hyperbolic itself. The theorem below generalizes this result.

Theorem 4.16. *Let R be a strongly relatively quasi–convex subgroup of G. Then R is a hyperbolic group.*

Proof. By Theorem 4.13, R is generated by a finite set Y. As in the proof of Theorem 4.13, for any $y \in Y$, let W_y denote a word over $X\cup\mathcal{H}$ representing y in G and let μ be defined by (4.5). We denote by $\Gamma(R, Y)$ the Cayley graph of R with respect to Y and define the map

$$\psi : \Gamma(R, Y) \to \Gamma(G, X\cup\mathcal{H})$$

as follows. First we require the restriction of ψ to the vertex set of $\Gamma(R, Y)$ to coincide with the natural embedding $R \to G$. Secondary, given an edge e of $\Gamma(R, Y)$ with label $\phi(e) = y$, we assume $\psi(e)$ to be the path in $\Gamma(G, X\cup\mathcal{H})$ labelled W_y. Obviously this two conditions uniquely define ψ. Note that ψ is a quasi–isometric embedding by Theorem 4.13. We denote by c_1, c_2 the corresponding quasi–isometry constants.

Let $\Delta = pqr$ be a geodesic triangle in $\Gamma(R, Y)$. Then the image $\Delta^* = p^*q^*r^*$ of Δ under ψ is a triangle in $\Gamma(G, X\cup\mathcal{H})$ whose sides are (λ, c)–quasi–geodesics, where λ and c depend only on c_1, c_2, not on Δ. Since $\Gamma(G, X \cup \mathcal{H})$ is a δ–hyperbolic space, by Lemma 3.5, there is a constant $H = H(\lambda, c, 0)$ such that each side of Δ^* belongs to the closed H–neighborhood (with respect to $dist_{X\cup\mathcal{H}}$) of the other two sides.

Let u be a vertex on one of the sides of Δ, say $u \in p$, and $u^* = \psi(u)$. Let also v^* be a vertex on $q^* \cup r^*$ such that

$$dist_{X\cup\mathcal{H}}(u^*, v^*) \leq H. \tag{4.6}$$

Clearly there exists a vertex $w^* \in q^* \cup r^*$ such that $w^* = \psi(w)$ for some $w \in q \cup r$ and

$$dist_{X\cup\mathcal{H}}(w^*, v^*) \leq \frac{1}{2}\mu. \tag{4.7}$$

Combining (4.6) and (4.7), we obtain

$$dist_Y(u, w) \leq c_1 dist_{X\cup\mathcal{H}}(u^*, w^*) + c_2 \leq c_1 \left(H + \frac{1}{2}\mu\right) + c_2.$$

Thus $\Gamma(R, Y)$ is δ'–hyperbolic for $\delta' = c_1 (H + 1/2\mu) + c_2$. □

Remark 4.17. The above theorem does not hold without the assumption $card\,(R \cap H_i^g) < \infty$. Indeed, let $G = H_1 * H_2$ for some finitely generated groups H_1, H_2. Let K_1 and K_2 be subgroups of H_1 and H_2 respectively. Suppose, in addition, that at least one of the subgroups K_1, K_2 is not finitely generated. Then the subgroup R generated by K_1, K_2 is obviously relatively quasi–convex, but not finitely generated by the Grushko–Neumann theorem since $R \cong K_1 * K_2$.

We conclude with a proposition describing the intersections of relatively quasi–convex subgroups. The logical scheme of the proof is due to Short [78].

Proposition 4.18. *Let P and R be two relatively quasi–convex subgroups of G. Then $P \cap R$ is relatively quasi–convex.*

Proof. Let us take an element $g \in P \cap R$ and consider a geodesic p in $\Gamma(G, X \cup \mathcal{H})$ such that $p_- = 1$, $p_+ = g$. We also take an arbitrary vertex $u \in p$. Let σ denote the quasi–convexity constant for both the subgroups P and R.

We have to show that there exists a vertex $v \in P \cap R$ within an X–distance at most σ' from v, where σ' is the constant which is independent of p and u. Let us consider a vertex $v \in P \cap R$ satisfying the following two conditions.

1. There exists a path s from u to v such that for every vertex $w \in s$ we have

   $$\max\{dist_X(w, P),\ dist_X(w, R)\} \leq \sigma.$$

   (Note that the set of vertices v satisfying this condition is non–empty; in particular, it contains g.)



Figure 4.3:

2. v is the closest vertex to u with respect to $dist_{X\cup\mathcal{H}}$ satisfying the first condition.

We are going to prove that

$$dist_{X\cup\mathcal{H}}(u,v) \le \left(card\ B_\sigma^X\right)^2, \tag{4.8}$$

where B_σ^X is the ball defined by (4.2).

Suppose that (4.8) is false. Then $l(s) > \left(card\ B_\sigma^X\right)^2$. According to the first condition there exist two vertices, say a and b, on s and two elements from B_σ^X, say x and y, such that $ax \in P$, $ay \in R$ and $bx \in P$, $by \in R$ (see Fig. 4.3). Let $s = s_1[a,b]s_2$. Consider the path t in $\Gamma(G, X\cup\mathcal{H})$ such that $t_- = u$ and $\phi(t) \equiv \phi(s_1)\phi(s_2)$. We state that $t_+ \in P\cap R$. Indeed $bx \in P$ and $b\overline{\phi(s_2)} = v \in P$ yield $x^{-1}\overline{\phi(s_2)} \in P$. Therefore,

$$t_+ = \overline{\phi(s_1)\phi(s_2)} = (ax)(x^{-1}\overline{\phi(s_2)}) \in P.$$

Similarly, $t_+ \in R$. Thus $t_+ \in P\cap R$. Moreover, for every vertex $w \in t$ we have

$$\max\{dist_X(w,P),\ dist_X(w,R)\} \le \sigma. \tag{4.9}$$

Indeed let $t = s_1t_2$, $\phi(t_2) \equiv \phi(s_2)$. If $w \in s_1$, then the fulfilment of (4.9) is obvious. Suppose that $w \in t_2$. Let z be the vertex of s_2 such that the segment

$[w, t_+]$ of t_2 has the same label as the segment $[z, v]$ of s_2. Then there exists $z' \in P$ such that $dist_X(z, z') \le \sigma$. Note that

$$\begin{aligned} w(z^{-1}z') = \quad & w(z^{-1}v)(v^{-1}z') = w\overline{\phi([z,v])}v^{-1}z' = w\overline{\phi([w,t_+])}v^{-1}z' = \\ & ww^{-1}t_+v^{-1}z' = t_+v^{-1}z' \in P \end{aligned}$$

as $t_+, v, z' \in P$. Thus

$$dist_X(w, P) \le dist_X(w, w(z^{-1}z')) = dist_X(z, z') \le \sigma.$$

Similarly $dist_X(w, R) \le \sigma$. We have proved that t_+ satisfies the first condition for v. Since t is shorter than s, we arrive at a contradiction.

Now we want to estimate the X–distance between u and v. Let r be an element of R such that

$$dist_X(u, r) \le \sigma. \tag{4.10}$$

Note that

$$dist_{X\cup\mathcal{H}}(v, r) \le dist_{X\cup\mathcal{H}}(u, v) + dist_{X\cup\mathcal{H}}(u, r) \le \sigma + \left(card\ B_\sigma^X\right)^2.$$

Let Y be a finite generating set for R. Then, according to Theorem 4.13, we have

$$dist_Y(v, r) \le c_1\left(\sigma + \left(card\ B_\sigma^X\right)^2\right) + c_2$$

where c_1, c_2 depend on R and G only. Therefore,

$$dist_X(v, r) \le dist_Y(v, r) \max_{y\in Y} |y|_X \le \left(c_1\left(\sigma + \left(card\ B_\sigma^X\right)^2\right) + c_2\right) \max_{y\in Y} |y|_X. \tag{4.11}$$

Summing (4.10) and (4.11), we obtain

$$dist_X(u, v) \le \left(c_1\left(\sigma + \left(card\ B_\sigma^X\right)^2\right) + c_2\right) \max_{y\in Y} |y|_X + \sigma.$$

Thus $P \cap R$ is relatively σ'–quasi–convex for

$$\sigma' = \left(c_1\left(\sigma + \left(card\ B_\sigma^X\right)^2\right) + c_2\right) \max_{y\in Y} |y|_X + \sigma.$$

□

4.3 Cyclic subgroups and translation numbers

It is well known that if a group G is hyperbolic, then any cyclic subgroup of G is quasi-isometrically embedded into G [45, 44]. This result is one of the corner stones of the small cancellation theory over hyperbolic groups [65].

It seems to be easy to prove the following by using arguments similar to the ordinary hyperbolic case. Let G be a group, $\{H_1, \dots, H_m\}$ a collection



Figure 4.4:

of subgroups of G. Suppose that the relative Cayley graph $\Gamma(G, X \cup \mathcal{H})$ is hyperbolic. Then for any element g that is not conjugate to an element of one of the subgroups $H_1, \ldots, H_m$, the cyclic subgroup generated by g is quasi–isometrically embedded into G (with respect to the relative metric on G).

Unfortunately, in general, this is not true. Indeed, consider an arbitrary groups H and the direct product $G = H_1 \times H_2$, where $H_i \cong H$, $i = 1, 2$. Then the relative Cayley graph $\Gamma(G, \mathcal{H})$ of G with respect to $\{H_1, H_2\}$ has finite diameter and, in particular, it is hyperbolic. To each element $h \in H$, one can assign the element $g = (h, h) \in H \times H$. Obviously g is not conjugate to an element of one of the copies of H in G whenever $h \neq 1$. However, the relative length of g^n is at most 2 for every n.

In this section we establish the quasi–convexity of cyclic subgroups generated by hyperbolic elements in relatively hyperbolic groups. As the previous example shows, to this end we need some additional arguments apart from the hyperbolicity of $\Gamma(G, X \cup \mathcal{H})$. Moreover, we do not restrict ourselves to the case of finitely generated groups, as the general case is important for the development of the small cancellation theory over relatively hyperbolic groups [68] and some of its applications. However the Splitting Theorem allows to reduce the proof to the finitely generated case. After such a reduction, we will follow the logical scheme suggested in [43] (the underlying idea has also been used in [35] and [3] to obtain similar results in wider contexts). Our main tools will be the theorems about quasi–convex subgroups and results from Section 3.4.

Theorem 4.19. *Let G be a finitely generated group hyperbolic relative to a collection of subgroups $\{H_1, \ldots, H_m\}$, g a hyperbolic element of G. Then the centralizer $C(g)$ of g in G is a strongly relatively quasi–convex subgroup in G.*

Proof. Let a be an element of $C(g)$, p a geodesic path in $\Gamma(G, X \cup \mathcal{H})$ such that $p_- = 1$, $p_+ = a$. We have to show that for any vertex v on p there exists a vertex $u \in C(g)$ such that $dist_X(u, v) \leq \sigma$, where $\sigma = \sigma(g)$ is independent of v and a. For this purpose, we also consider a geodesic q in $\Gamma(G, X \cup \mathcal{H})$ such that

$\phi(p) \equiv \phi(q)$ and $q_- = g$ (see Fig. 4.4). Since $a \in C(g)$, we have

$$q_+ = g\overline{\phi(q)} = g\overline{\phi(p)} = ga = ag.$$

By k we denote the length $|g|_X$. Then,

$$dist_X(p_+, q_+) = dist_X(a, ag) = |g|_X.$$

Thus (p, q) is a pair of k–similar symmetric geodesics in $\Gamma(G, X \cup \mathcal{H})$.

Let $\varkappa = \varkappa(k)$ be the constant from Theorem 3.37, w the vertex on q synchronous to v. By Theorem 3.37, $dist_X(v, w) \leq \varkappa$. Let $Conj$ denote the set of all pair of elements $(f, g) \in G \times G$ such that f and g are conjugate in G. To each such a pair we assign an element $t_{f,g} \in G$ such that $f^{t_{f,g}} = g$ and set

$$\sigma = \max\{|t_{f,g}|_X \ : \ (f, g) \in Conj, \ |f|_X + |g|_X \leq k + \varkappa\}.$$

Since G is locally finite with respect to the metric $dist_X$, σ is well–defined and depends on k and $\varkappa$ only (the collection of elements $t_{f,g}$ is supposed to be fixed).

By the choice of σ, there exists an element $t \in G$ of X–length at most σ such that $(v^{-1}w)^t = g$. Note also that $w = g\overline{\phi([g, w])} = gv$. This yields

$$vtg = vt(v^{-1}w)^t = v(v^{-1}w)t = wt = gvt.$$

Hence $vt \in C(g)$. It remains to note that $dist_X(v, vt) = |t|_X \leq \sigma$.

Let us show that $C(g) \cap H_i^f$ is finite for any $f \in G$, $i = 1, \ldots m$. Evidently we have $C(g) \cap H_i^f \leq H_i^{fg}$. Therefore,

$$C(g) \cap H_i^f \leq H_i^{fg} \cap H_i^f = \left(H_i^{fgf^{-1}} \cap H_i\right)^f.$$

Since g is hyperbolic, $fgf^{-1} \notin H_i$ and thus the intersection $H_i^{fgf^{-1}} \cap H_i$ is finite by Proposition 2.36. The lemma is proved. □

Corollary 4.20. *Let G be a finitely generated group hyperbolic relative to a collection of subgroups $\{H_1, \ldots, H_m\}$, $g \in G$ a hyperbolic element of infinite order. Then there exist $\lambda > 0, c \geq 0$ such that*

$$|g^n|_{X \cup H} \geq \lambda|n| - c \tag{4.12}$$

for any $n \in \mathbb{Z}$.

Proof. By Lemma 4.19, $C(g)$ is strongly relatively quasi–convex. Further according to Theorems 4.13 and 4.16, $C(g)$ is generated by a finite set Y and hyperbolic. The center Z of $C(g)$ is infinite, as it contains $\langle g \rangle$. As is well–known, any hyperbolic group with infinite center is virtually cyclic. Hence the index of $\langle g \rangle$ in $C(g)$ is finite. This obviously implies that the map $(\langle g \rangle, dist) \to (C(g), dist_Y)$ is a quasi–isometric embedding (here $dist$ denotes the natural metric on $\langle g \rangle$ with respect to the generating set $\{g\}$).

Thus we have a sequence of quasi–isometric embeddings

$$(\langle g\rangle, dist) \to (C(g), dist_Y) \to (G, dist_{X\cup\mathcal{H}}).$$

Since composition is also a quasi–isometric embedding, we obtain (4.12). □

Corollary 4.21. *Let g be a hyperbolic element of infinite order in G. If $(g^k)^t = g^l$ for some $k, l \in \mathbb{Z}$, $t \in G$, then $k = \pm l$.*

Proof. The argument is standard. If $k \neq \pm l$, we can assume $|k| < |l|$. Then $(g^{k^n})^{t^n} = g^{l^n}$. This yields

$$|g^{l^n}|_{X\cup\mathcal{H}} = \quad |(g^{k^n})^{t^n}|_{X\cup\mathcal{H}} \leq 2n|t|_{X\cup\mathcal{H}} + |k|^n|g|_{X\cup\mathcal{H}} =$$

$$2n|t|_{X\cup\mathcal{H}} + (l^n)^{\log_l k}|g|_{X\cup\mathcal{H}}$$

for any $n \in \mathbb{N}$ contradictory (4.12). □

The reader can also derive

Corollary 4.22. *Suppose that B is a subgroup of G. If B is isomorphic to a Baumslag–Solitar group, that is,*

$$B \cong \langle a, b \mid (a^k)^b = a^l\rangle,$$

then B is conjugate to a subgroup of H_λ for some $\lambda \in \Lambda$.

Our next goal is to show that, in fact, the constant λ in Corollary 4.20 is independent of g. It is convenient to express this property in terms of translation numbers.

Definition 4.23. Let K be a group generated by a finite set S relative to a collection of subgroups $\{I_\lambda\}_{\lambda\in\Lambda}$, x an element of K. The *relative translation number* of x is defined to be

$$\tau^{rel}(x) = \lim_{n\to\infty} \frac{1}{n}|x^n|_{S\cup\mathcal{I}}, \tag{4.13}$$

where $|\cdot|_{S\cup\mathcal{I}}$ denotes the relative length function with respect to S and $\{I_\lambda\}_{\lambda\in\Lambda}$.

The following lemma is quite trivial (The proof of its non–relative analog can be found in [43]; it works in the relative case without any changes.)

Lemma 4.24. *1. The limit in (4.13) always exists.*

2. $\tau^{rel}(x) = \inf\limits_{n} \frac{|x|_{S\cup\mathcal{I}}}{n}$. In particular, $\tau^{rel}(x) \leq |x|_{S\cup\mathcal{I}}$.

3. $\tau^{rel}(x) = \tau^{rel}(x^t)$ for any $x, t \in K$; thus $\tau^r el(x)$ depends only on the conjugacy class of x.

4. $\tau^{rel}(x^n) = |n|\tau^{rel}(x)$ for any $x \in K$, $n \in \mathbb{Z}$.

Theorem 4.25. *Let G be a group, $\{H_\lambda\}_{\lambda\in\Lambda}$ a collection of subgroups of G. Suppose that G is hyperbolic with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$. Then there exists $d > 0$ such that for every hyperbolic element of infinite order $g \in G$, the relative translation number of g with respect to $\{H_\lambda\}_{\lambda\in\Lambda}$ satisfies the inequality $\tau^{rel}(g) > d$.*

Proof. The proof consists of several lemmas. First of all we are going to reduce the proof to the case when G is finitely generated.

Lemma 4.26. *Let K be the fundamental of a tree of groups with vertex groups $I_1, \ldots, I_m$. Suppose that x is an element of K. Then either x is conjugate to an element of I_i for some i or the translation number of g relative to $I_1, \ldots, I_m$ is at least 2.*

Proof. Obviously it suffices to prove the lemma in case $m = 2$. (Then we can apply inductive arguments). If $m = 2$, K is a free product of I_1, I_2 with amalgamated subgroups, say A and B. Below we use terminology and some well–known results about amalgamated products, which can be found, for example, in [58, Sec. 4.2]. If x is not conjugate to an element of I_1 or I_2, then x is conjugate to a cyclically reduced (in the sense of amalgamated products) element $y = ay_1 \ldots y_d$, where $d \geq 2$, $a \in A$, $y_1, \ldots, y_d$ are coset representatives of I_1 or I_2 with respect to A, and y_i, y_{i+1} (indices are *mod* d) are not in the same factor. Then the number of factors in the reduced form of y^n is at least $2dn \geq 2n$. Since the reduced form is unique and the reduction process does not increase the number of factors, we have $|y^n|_{\mathcal{I}} \geq 2|n|$. This yields the assertion of the lemma as translation numbers depend only on conjugacy classes. $\square$

Recall that by the Splitting Theorem, $G = G_0 * \left(*_{\lambda\in\Lambda\setminus\Lambda_0} H_\lambda\right)$, where G_0 is a tree of groups with vertex groups $H_\lambda, \lambda \in \Lambda_0 = \{\lambda_1, \ldots, \lambda_m\}$, and Q. If $g \in G$ is hyperbolic, then either g has length at least 2 (as the element of a free product), or g is conjugate to an element of G_0. In the first case $\tau^{rel}(g) \geq 2$ by the obvious reasons. In the second case Lemma 4.26 shows that it suffices to prove the theorem for elements g such that g is conjugate to an element of Q. Since translation numbers depend only on the conjugacy classes, we may assume $g \in Q$. Recall that Q is finitely generated and hyperbolic relative to some subgroups L_i, $i = 1, \ldots, m$. By Proposition 2.49, it suffices to show that there exists $d > 0$ such that for any $g \in Q$ the relative translation number of g in Q with respect to $\{L_1, \ldots, L_m\}$ is greater than d.

Thus we can assume that the group G is generated by a finite set X in the usual (non–relative) sense. To prove the theorem it suffices to show that for some $a > 0$, the set

$$T_G(a) = \{\tau^{rel}(g) \mid g \in G,\ g \text{ is hyperbolic},\ \tau^{rel}(g) < a\} \tag{4.14}$$

is finite. To this end we use an auxiliary lemma below, which can be regarded as a generalization of Lemma 4.5. Throughout the rest of this section, we use the notation M, L (see the beginning of the previous chapter) and denote by δ the hyperbolicity constant of $\Gamma(G, X \cup \mathcal{H})$.

Lemma 4.27. *Suppose that g is a hyperbolic element of G satisfying the following conditions.*

1. *g has minimal relative length among all elements of the conjugacy class g^G.*
2. *$|g|_{X\cup\mathcal{H}} \le 8\delta + 1$.*
3. *$\tau^{rel}(g) < \xi$ for $\xi = \frac{1}{8\delta+3}$.*

Then the X–length of G satisfies

$$|g|_X \le (8\delta + 1)(32\delta + 6)ML. \tag{4.15}$$

Proof. Since $\tau^{rel}(g) < \xi$, there exists $N \in \mathbb{N}$ such that $|g^N|_{X\cup\mathcal{H}} < \xi N$. Let $U \in (X \cup \mathcal{H})^*$ be a shortest word representing g in G and $W \in (X \cup \mathcal{H})^*$ a shortest word representing g^N. Obviously we have

$$\|U\| \le 8\delta + 1 \tag{4.16}$$

and

$$\|W\| \le \xi N. \tag{4.17}$$

We consider the cycle pq^{-1} in $\Gamma(G, X \cup \mathcal{H})$ such that $p_- = q_- = 1$, $\phi(p) \equiv U^N$, and $\phi(q) \equiv W$. There are three possibilities to consider.

Case 1. First suppose that any component of pq^{-1} is isolated. Given an H_i–syllable V of U, we have at least N H_i–components of p labelled V. By Lemma 3.1, we have

$$N|\overline{V}|_X \le MLl(pq^{-1}) \le ML(N\|U\| + \|W\|).$$

Using (4.16), (4.17) and dividing both the sides of the previous inequality by N, we obtain

$$|\overline{V}|_X \le ML\frac{N(8\delta + 1) + \xi N}{N} < ML(8\delta + 2) \tag{4.18}$$

(note that $\xi < 1$). Since (4.18) is true for every syllable of U, we obtain

$$|g|_X = |\overline{U}|_X \le \|U\|ML(8\delta + 2) \le (8\delta + 1)(8\delta + 2)ML.$$

Obviously this inequality is even stronger than (4.15).

Case 2. Assume that there are two connected components s_1 and s_2 of p. Then repeating the same arguments as in the proof of Lemma 4.5, we obtain

$$|g|_X \le 2ML(8\delta + 1)$$

and the inequality (4.16) obviously holds. We leave details to the reader.

Case 3. Suppose that no different components of p are connected. As the path q is geodesic, no different components of q are connected. Therefore, the only possibility for two components s and t of pq^{-1} to be connected is $s \in p$, $t \in q$.

Let $s_1, \ldots, s_n$ be the set of all components of p such that for any $i = 1, \ldots, n$, there exists a component of q, denoted by t_i, that is connected to s_i. Let us denote by e_i and f_i the paths of length at most 1 in $\Gamma(G, X \cup \mathcal{H})$ such that $(e_i)_- = (t_i)_-$, $(e_i)_+ = (s_i)_-$, $(f_i)_- = (t_i)_+$, $(f_i)_+ = (s_i)_+$. We also set e_0, f_0 (respectively e_{n+1} and f_{n+1}) to be the paths consisting of just one vertex 1 (respectively g^N). Without loss of generality we may assume that

$$p = r_1 s_1 \ldots r_n s_n r_{n+1}.$$

Let us choose a subsequence $s_{i_1}, \ldots, s_{i_l}$ of the sequence $s_1, \ldots, s_n$ as follows. We set $s_{i_1} = s_1$. Further suppose we have already chosen s_{i_k}. Then $s_{i_{k+1}}$ is defined to be the first component in the sequence $s_{i_k+1}, s_{i_k+2}, \ldots, s_n$ such that the corresponding component $t_{i_{k+1}}$ belongs to the segment $[(t_{i_k})_+, q_+]$ of q. Thus after completing this process, we will obtain a sequence of components $s_{i_1}, \ldots, s_{i_l}$ of p such that (see Fig. 4.5):

(i) $p = p_1 s_{i_1} \ldots p_l s_{i_l} p_{l+1}$ for some $p_1, \ldots p_{l+1}$;

(ii) $q = q_1 t_{i_1} \ldots q_l t_{i_l} q_{l+1}$ for some $q_1, \ldots, q_{l+1}$;

(iii)for any $j = 1, \ldots l+1$, every component of the cycle $c_j = f_{i_{j-1}} p_j e_{i_j}^{-1} q_{i_j}^{-1}$ is isolated in c_j (Figure 4.5).

For simplicity, we change the notation and denote s_{i_j}, f_{i_j}, and e_{i_j} by s_j, f_j, and e_j respectively. Thus we have

$$p = p_1 s_1 \ldots p_l s_l p_{l+1},$$

$$q = q_1 t_1 \ldots q_l t_l q_{l+1},$$

and

$$c_j = f_{j-1} p_j e_j^{-1} q_j^{-1}.$$

We will call $c_1, \ldots, c_{l+1}$ sections. We state that there exists a section c_j such that the following two inequalities hold.

$$\left\{ \begin{array}{l} l(q_j) \leq l(p_j), \\ \\ l(p_j) \geq 8\delta + 1. \end{array} \right. \tag{4.19}$$

Indeed let us denote by S_1 the set of all sections c_j such that $l(q_j) > l(p_j)$ and by S_2 the set of all sections c_j such that $l(p_j) < 8\delta + 1$. It suffices to show that there exists a section $c_j \notin S_1 \cup S_2$. Notice that the number of all sections is at most $l(q) < \xi N$. According to the choice of U, any component s_j consists



Figure 4.5:

of a single edge. Let $\pi(c_j)$ denote the number $l(p_j) + l(s_j) = l(p_j) + 1$. On one hand, we have

$$\sum_{j=1}^{l+1} \pi(c_j) = l(p) + 1.$$

On the other hand,

$$\sum_{c \in S_2} \pi(c) < (8\delta + 2) card\, S_2 < (8\delta + 2)\xi N \tag{4.20}$$

and

$$\sum_{c \in S_1} \pi(c) < l(q) + 1 < \xi N + 1 \tag{4.21}$$

Inequalities (4.20) and (4.21) yield

$$\sum_{c \in S_1 \cup S_2} \pi(c) < (8\delta + 3)\xi N + 1 = N + 1 \leq l(p) + 1.$$

Therefore there is a section which is not in $S_1 \cup S_2$.

Let c_j be a section satisfying (4.19). We have

$$\phi(p_j) \equiv U_0^k A,$$

where U_0 is a cyclic shift of the word U and $\|A\| < \|U\|$. According to the second inequality of (4.19) and the second condition of the lemma, we have $k \geq 1$. Let V be a syllable of U. Then arguing as in Case 1, and taking into account the inequality

$$l(c_j) \leq l(p_j) + l(q_j) + 2 \leq 2l(p_j) + 2 < 2(k+1)\|U\| + 2 \leq 2(k+1)(8\delta + 1) + 2$$

we obtain

$$k|\overline{V}|_X \leq MLl(c_j) \leq ML(2(k+1)(8\delta+1)+2).$$

Hence

$$|\overline{V}|_X \leq ML(32\delta+6).$$

Since this is true for every syllable of U, we obtain

$$|g|_X = |\overline{U}|_X \leq \|U\| ML(32\delta+6) \leq (8\delta+1)(32\delta+6)ML.$$

□

Let us return to the proof of Theorem 4.25. Given an arbitrary element g in G which has shortest relative length in the conjugacy class g^G, there are two possibilities.

First suppose that $|g|_{X\cup\mathcal{H}} \geq 8\delta+1$. We take a shortest word $U \in (X\cup\mathcal{H})^*$ representing g and consider the path p_n such that $(p_n)_- = 1$, $\phi(p_n) \equiv U^n$, $n \in \mathbb{N}$. Since g is a shortest element in g^G, p_n is $(8\delta+1)$–local geodesic in $\Gamma(G, X\cup\mathcal{H})$. Therefore, by Lemma 4.3 we have

$$|g^n|_{X\cup\mathcal{H}} = dist_{X\cup\mathcal{H}}(1, (p_n)_+) \geq \frac{1}{\lambda}(l(p_n)-c) = \frac{1}{\lambda}(n|g|_{X\cup\mathcal{H}}-c),$$

for $\lambda \leq 3$ and $c = 2\delta$. Hence,

$$\tau^{rel}(g) \geq \lim_{n\to\infty} \frac{1/\lambda(n|g|_{X\cup\mathcal{H}}-c)}{n} = \frac{1}{\lambda}|g|_{X\cup\mathcal{H}} \geq 1/3.$$

Now assume that $|g|_{X\cup\mathcal{H}} < 8\delta+1$. Then, by Lemma 4.27, either $\tau^{rel}(g) \geq \xi = 1/(8\delta+3)$ or g is conjugate to an element of the set $B = B^X_{(8\delta+1)(32\delta+6)ML}$ of elements whose X–length is at most $(8\delta+1)(32\delta+6)ML$. Since B is finite and $\tau^{rel}(g)$ depends only on the conjugacy class of g, $T_G(a)$ is finite for $a = 1/(8\delta+3)$. □

Chapter 5

Algorithmic problems

5.1 The word and membership problems

Recall that the word problem for a recursively presented group generated by a recursive set X is to decide, given a word W in the alphabet $X^{\pm 1}$, whether W represents 1 in G. In [37], Farb showed that the word problem is solvable for any finitely generated group G hyperbolic relative to subgroups $\{H_1, \ldots, H_m\}$ provided it is solvable for each of the subgroups $H_1, \ldots, H_m$. It is not hard to generalize this result as follows.

Theorem 5.1. *Suppose that G is a group given by a finite relative presentation with respect to recursively presented subgroups $H_1, \ldots, H_m$. Assume also that the corresponding relative Dehn function $\delta^{rel}(n)$ is bounded from above by some recursive function and the word problem is solvable for all subgroups H_i, $i = 1, \ldots, m$. Then the word problem is solvable for G.*

Proof. Let X be a finite generating set of G, $X = X^{-1}$, $\widetilde{H}_i$ an isomorphic copy of H_i, $i = 1, \ldots, m$,

$$F = F(X) * \widetilde{H}_1 * \ldots * \widetilde{H}_m,$$

and let N be the kernel of the naturally defined homomorphism $F \to G$. Suppose that $N = \langle \mathcal{R} \rangle^F$ for a certain finite subset $\mathcal{R} \subset N$, and the corresponding relative Dehn function $\delta^{rel}(n)$ is bounded from above by some recursive function. We also assume that the relative presentation

$$\langle X, H_1, \ldots, H_m \mid R = 1,\ R \in \mathcal{R} \rangle \tag{5.1}$$

is reduced (see Definition 2.24). Let Ω_i, Ω denote the (finite) sets introduced in Definition 2.25. By Proposition 2.29, Ω_i generates H_i. Thus the group F is generated by the finite set $Z = X \cup \widetilde{\Omega}$, where $\widetilde{\Omega}$ consists of the preimages of elements of Ω under the canonical homomorphisms $\widetilde{H}_i \to H_i$.

For any word W over X, $W \in N$, of length $\|W\| \le n$, we consider a van Kampen diagram Δ of minimal type over (5.1) with boundary label W. Note that any edge of this diagram is labelled by an element of Z. Indeed this is so for any external edge since W is a word over X. If e is an internal edge, then e belongs to the boundary of some $\mathcal{R}$–cell by Lemma 2.15 and thus $\phi(e) \in Z$. Fixing a basepoint in Δ, we obtain a representation

$$W =_F \prod_{i=1}^{k} f_i^{-1} R_i f_i, \tag{5.2}$$

$f_i \in F$, $R_i \in \mathcal{R}$, $i = 1, \ldots, k$, where $k \le \delta^{rel}(n)$, and each element f_i is a label of a path without self–intersections in Δ. In particular, $|f_i|_Z$ is not greater than the number of all edges in Δ, i.e.,

$$|f_i|_Z \le M\delta^{rel}(n) + \|W\| \le M\delta^{rel}(n) + n$$

(recall that $M = \max\limits_{R\in\mathcal{R}} \|R\|$.) Thus the number of factors in (5.2) and the lengths of conjugating elements f_i with respect to the (finite) set Z are bounded by recursive functions of n. Since the word problem is solvable for F, these bounds allow to derive that it is solvable for G. $\square$

It is worth to notice that one can easily provide an example of a finitely generated group G and a finitely generated subgroup H of G such that the word problem is solvable for both G and H, but the corresponding relative Dehn function is not well–defined. (For example, this is so if $G = H \times \mathbb{Z}$ and $H = \mathbb{Z}$.)

Recall that the *membership* problem for a subgroup K of a group G is to decide for a given element $g \in G$ whether g belongs to K.

Theorem 5.2. *Suppose that G is a group given by a finite relative presentation with respect to recursively presented subgroups $H_1, \ldots, H_m$. Assume also that the corresponding relative Dehn function $\delta^{rel}(n)$ is bounded from above by some recursive function and the word problem is solvable for all subgroups H_i, $i = 1, \ldots, m$. Then the membership problem is solvable for H_i, $i = 1, \ldots, m$.*

Note that the theorem does not hold without the requirement of the solvability of the word problem in H_i, $i = 1, \ldots, m$. For example, if H is a finitely presented group with undecidable word problem, then the group $G = H$ is hyperbolic relative to the subgroups $H_1 = H$ and $H_2 = \{1\}$. Clearly the membership problem for H_2 is unsolvable in this case. To prove the theorem we need an auxiliary definition.

Definition 5.3. Let H be a group generated by a finite set A and K a subgroup of H generated by a finite set B. The *distortion function* of K in H with respect to the generating sets A and B is defined to be

$$\Delta_K^H(n) = \max\{|x|_B \mid x \text{ satisfies } |x|_A \le n\}.$$

It is easy to see that Δ_K^H is independent (up to equivalence) of the choice of finite generating sets in H and K. In case $\Delta_K^H(n) \sim n$, we say that K is undistorted in H.

Lemma 5.4. *Let G be a finitely generated group, $\{H_1, \dots H_m\}$ a collection of finitely generated subgroups of G. Suppose that G is finitely presented with respect to $\{H_1, \dots H_m\}$ and the corresponding relative Dehn function $\delta^{rel}(n)$ is well–defined. Then for any $i = 1, \dots, m$, the distortion of the subgroup H_i in G satisfies*

$$\Delta^G_{H_i}(n) \preceq \delta^{rel}(n). \tag{5.3}$$

Proof. Let h be a non–trivial element of H_i, W a shortest word over X representing h. Then there exists a cycle $c = pq^{-1}$ in $\Gamma(G, X \cup \mathcal{H})$ such that $\phi(p) \equiv W$ and q is an edge in $\Gamma(G, X \cup \mathcal{H})$ labelled h. Note that q is an isolated H_i–component of c. Applying Lemma 2.27 we obtain

$$|h|_{\Omega_i} = |\overline{\phi(q)}|_{\Omega_i} \leq M Area^{rel}(c) \leq M\delta^{rel}(\|W\| + 1) = M\delta^{rel}(|h|_X + 1).$$

This implies (5.3). □

Proof of Theorem 5.2. As is well-known [36], if the word problem in a finitely generated group H is solvable and a subgroup $K \leq H$ is finitely generated, then the membership problem for K is solvable if and only if $\Delta^H_K(n)$ is bounded from above by a recursive function. Thus Theorem 5.2 follows from the previous lemma. □

Corollary 5.5. *Let G be a finitely generated group hyperbolic relative to recursively presented subgroups $\{H_1, \dots, H_m\}$. Suppose that the word problem is solvable in each of the subgroups $H_1, \dots, H_m$. Then:*

1) (Farb, [37]) The word problem is solvable in G.

2) For any $i = 1, \dots, m$, the membership problem is solvable for H_i.

5.2 The parabolicity problems

In the previous chapter we saw that some important properties (such as the finiteness of the conjugacy classes of elements of finite orders, strong quasi–convexity of cyclic subgroups, etc.) hold for hyperbolic elements although they can be violated for parabolic ones. Thus given a finitely generated group G which is hyperbolic relative to a collection of subgroups $\{H_1, \dots, H_m\}$, it is natural to consider the following two algorithmic problems

1) (*The general parabolicity problem*) Given an element $g \in G$, decide whether g is parabolic or hyperbolic.

2) (*The special parabolicity problem*) Given an element $g \in G$ and $i \in \{1, \dots, m\}$, decide whether g is conjugate to an element of H_i.

In case $m = 1$ these problems coincide. It is proved in [20] that they are solvable for any finitely generated group G hyperbolic relative to a subgroup H whenever the conjugacy problem is decidable in H. Similar arguments allow to show that the special parabolicity problem is solvable for any finitely generated group G hyperbolic relative to $\{H_1, \ldots, H_m\}$ whenever the conjugacy problem is solvable in all $H_1, \ldots, H_m$. We observe that the last requirement is essential for the special parabolicity problem. On the other hand, the general parabolicity problem is solvable in G whenever the word problem is solvable in $H_1, \ldots, H_m$. In this section we give the proof of these results in the spirit of our paper.

Theorem 5.6. *Let G be a group hyperbolic relative to a collection of recursively presented subgroups $\{H_1, \ldots, H_m\}$.*

1) If the word problem is solvable for all H_i, $i = 1, \ldots, m$, then the general parabolicity problem is solvable in G. Moreover, there is an algorithm which allows, given a parabolic element $g \in G$, to find some $t \in G$ and some $j \in \{1, \ldots, m\}$ such that $g^t \in H_j$.

2) (Bumagin, [20]) If the conjugacy problem is solvable for all H_i, $i = 1, \ldots, m$, then the special parabolicity problem is solvable in G. Moreover, there is an algorithm which allows, given $i \in \{1, \ldots, m\}$ and $g \in G$ that is conjugate to an element of H_i, to find an element $t \in G$ such that $g^t \in H_i$.

The proof of the theorem is based on the next two lemmas.

Lemma 5.7. *Let G be a finitely generated group hyperbolic relative to a collection of subgroups $\{H_1, \ldots, H_m\}$. There exists a recursive function $\sigma(k)$ satisfying the following condition. Let g be a parabolic element of G such that $|g|_X \leq k$. Then there exists $t \in G$ such that $g^t \in H_j$ for a certain $j \in \{1, \ldots, m\}$ and $|t|_X \leq \sigma(k)$.*

Proof. Let $\mathcal{P}$ be the set of all pairs of symmetric geodesics in $\Gamma(G, X \cup \mathcal{H})$ with characteristic elements g, h, where $h \in H_j$ for some $j = 1, \ldots, n$ (see Section 3.4 for definitions). Let (p, q) be a pair of geodesics of minimal lengths in $\mathcal{P}$. By Corollary 3.42, the length of the element $t = \overline{\phi(p)} = \overline{\phi(q)}$ is not greater than $\rho(k)$, where the constant $\rho(k)$ can be effectively calculated (one can notify that this is the common property of all constants in our paper). We note that no synchronous components of p and q are connected. Indeed if $p = p_1 a p_2$, $q = q_1 b q_2$, where a, b are connected synchronous components of p and q respectively, then $(p_1, q_1) \in \mathcal{P}$, which contradicts to the minimality of length of p and q. Therefore, by Lemma 3.39 there are no connected components of p and q at all.

Further let r be the edge in $\Gamma(G, X \cup \mathcal{H})$ labelled by an element $h \in H_j$ for some j such that $r_- = p_+$, $r_+ = q_+$. Note that r can not be connected to an H_j–component of p or q. Indeed if $p = s_1 c s_2$, where c is an H_j–component of p connected to r, then the label of cs_2 represents an element of H_j. Thus, for $t_0 = \overline{\phi(s_1)}$, we have

$$g^{t_0} = g^{t\overline{\phi(cs_2)}^{-1}} = h^{\overline{\phi(cs_2)}^{-1}} \in H_i.$$

This contradicts to the choice of (p, q) again. Hence no components of the paths pr and q are connected. Obviously pr and q are k–connected $(1, 2)$–quasi–geodesics. By Theorem 3.23, any component s of p has X–length at most

$|s|_X \leq \varepsilon(1,2,k)$, where $\varepsilon(1,2,k)$ can be effectively calculated for given k. Thus we have $|t|_X \leq \sigma(k)$ for $\sigma(k) = \rho(k)\varepsilon(1,2,k)$. □

Lemma 5.8. *Let G, $\{H_1, \ldots, H_m\}$ be as in the previous lemma. Suppose, in addition, that $H_1, \ldots, H_m$ are recursively presented and have solvable conjugacy problem. Then there exists a recursive function $\theta(k)$ satisfying the following condition. Let g be an element of G conjugate to an element $h \in H_i$ for some $i \in \{1, \ldots, m\}$ such that $|g|_X \leq k$. Then there exists $t \in G$ such that $g^t \in H_i$ and $|t|_X \leq \theta(k)$.*

Proof. Let $\mathcal{Q}$ be the set of all pairs of symmetric geodesics in $\Gamma(G, X \cup \mathcal{H})$ with characteristic elements g, h_0, where $h_0 \in H_i$. Let (p,q) be a pair of geodesics of minimal lengths in $\mathcal{Q}$. As above, using Corollary 3.42, we obtain $|t|_{X\cup\mathcal{H}} \leq \rho(k)$, where $t = \overline{\phi(p)} = \overline{\phi(q)}$. By Lemma 3.39, only synchronous components of p and q can be connected.

By r we denote the edge in $\Gamma(G, X \cup \mathcal{H})$ going from p_+ to q_+ and labelled by the element $h_0 \in H_i$, which is conjugate to g by t. The paths pr and q are k–similar $(1,2)$–quasi–geodesics. As in the previous lemma, we can show that r can not be connected to a component of p or q.

If there are no connected synchronous components of p and q, we can repeat the arguments from the proof of the previous lemma. Further let $p = p_1 a_1 \ldots p_n a_n p_{n+1}$, $q = q_1 b_1 \ldots q_n b_n q_{n+1}$, where for $j = 1, \ldots, n$, a_j, b_j are connected synchronous components of p and q respectively. We assume that p_j and q_j contain no connected components for $j = 1, \ldots, n+1$.

We denote by v_j and w_j the elements $((a_j)_-)^{-1}(b_j)_-$ and $((a_j)_+)^{-1}(b_j)_+$ of G respectively. By Theorem 3.23, for any $j = 1, \ldots, n$, we have $\max\{|v_j|_X, |w_j|_X\} \leq \varepsilon(1,2,k)$. Since the conjugacy problem is decidable in $H_1, \ldots, H_m$, there is a recursive function $\tau : \mathbb{N} \to \mathbb{N}$ such that for every $i \in \{1, \ldots, m\}$, and any two elements $v, w \in H_i$ that are conjugate in H_i, there is an element $s \in H_i$ such that $v^s = w$ and $|s|_X \leq \tau(\max\{|v|_X, |w|_X\})$. Note that elements v_j and w_j are conjugate in the subgroup H_{i_j} for a suitable i_j. Let $s_j \in H_{i_j}$ be the corresponding conjugating element such that $v_j^{s_j} = w_j$ and $|s_j|_X \leq \tau(\varepsilon(1,2,k))$. It is easy to check that for the element $z = \overline{\phi(p_1)} s_1 \ldots \overline{\phi(p_n)} s_n \overline{\phi(p_{n+1})}$, we have $g^z = g^t = h_0$.

Let us estimate the length of z. By our assumption, no components of subpaths p_j and q_j, $j = 1, \ldots, n$ are connected. The same is true for the subpaths $p_{n+1}r$ and q_n. Since for any $j = 1, \ldots, n$, p_j and q_j are $l = \max\{\varepsilon(1,2,k), k\}$–similar, as well as $p_{n+1}r$ and q_{n+1}, for any component a of p_j, $j = 1, \ldots, n+1$, we have $|a|_X \leq \varepsilon(1,2,l)$ by Theorem 3.23. Hence,

$$|t|_X = |\overline{\phi(p)}|_X \leq \rho(k) \max\{\varepsilon(1,2,l),\, \tau(\varepsilon(1,2,k))\}.$$

□

Proof of Theorem 5.6. Let us prove the first statement of the theorem. By Corollary 5.5, the word problem is decidable in G. Moreover, the membership problem is decidable for H_i, $i = 1, \ldots, m$. Given an element $g \in G$, we solve the membership problem for all elements of type g^t, where $|t|_X \le \sigma(|g|_X)$, and H_j for all $j = 1, \ldots, m$. By Lemma 5.7, g is parabolic if and only if $g^t \in H_j$ for some $|t|_X \le \sigma(|g|_X)$ and $j \in \{1, \ldots, m\}$.

The proof of the second statement follows from Lemma 5.8 in the same way. □

Finally we note that the solvability of the conjugacy problem in each H_i, $i = 1, \ldots, m$, is essential in the second statement of Theorem 5.6. That is, if we replace this condition with the solvability of the word problem in each H_i, $i = 1, \ldots, m$, the statement would be false. Clearly the minimal possible value of m in any counterexample is 2.

Theorem 5.9. *There exists a group G hyperbolic relative to finitely presented subgroups H_1, H_2 such that the word problem is solvable in H_1 and H_2 and the special parabolicity problem is unsolvable in G.*

Proof. Suppose that H is a finitely generated group having solvable word problem, $x \in H$ a fixed nontrivial element of order 2 such that there is no algorithm which allows to decide whether $h \sim x$ for a given element $h \in H$. (We explain how to construct such a group below.) Let $G = H_1 *_{\langle x \rangle} H_2$ be the amalgamated product of two copies of H, where the amalgamated subgroups are generated by the elements corresponding to x in each copy of H. Note that G is hyperbolic relative to $\{H_1, H_2\}$. Indeed the existence of the action of G on the Bass–Serre tree yields that G is hyperbolic relative to $\{H_1, H_2\}$ in the sense of Bowditch and thus in our sense (see Appendix). Obviously an element $h \in H_1$ is conjugate to an element of H_2 if and only if h is conjugate to x in H_1. Thus the special parabolicity problem is unsolvable in G.

To construct the group H with the desired properties, let us take the abelian group

$$A = \langle x_i, i \in \mathbb{N} \mid x_i^2 = 1,\, [x_i, x_j] = 1,\, i, j \in \mathbb{N} \rangle.$$

Let $f : \mathbb{N} \to \mathbb{N}$ be a recursive function such that the range of f is not recursive. Set $N = \{x_{f(i)},\, i \in \mathbb{N}\}$. We fix an arbitrary element $x \in N$ and consider the sequence of groups $Q(n)$, $n = 0, 1, \ldots$, such that $Q(0) = A$ and $Q(n)$ is obtained from $Q(n-1)$ by adding one extra generator t_n subject to the relation

$$t_n^{-1} x_{f(n)} t_n = x.$$

Lemma 5.10. *For any non–negative integer n, the word problem and in $Q(n)$ is decidable.*

Proof. The case $n = 0$ is obvious. Let $n > 0$. The group $Q(n)$ is the HNN–extension of $Q(n-1)$ with finite associated subgroups $\langle x_{f(n)} \rangle$ and $\langle x \rangle$. Since the subgroups $\langle x_{f(n)} \rangle$ and $\langle x \rangle$ are finite and the word problem in $Q(n-1)$ is solvable, the membership problem for $\langle x_{f(n)} \rangle$ and $\langle x \rangle$ is solvable in $Q(n-1)$. Hence the word problem is solvable in $Q(n)$ [56, Corollary 2.2, Ch. IV]. □

We set $Q = \bigcup_{i=1}^{\infty} Q(n)$. Clearly Q is recursively presented and the word problem is solvable in Q.

Lemma 5.11. *For an element $a \in A$, a is conjugate to x in Q if and only if $a \in N$.*

Proof. The 'if' part follows from our construction. We now suppose that $a \in A$ and a is conjugate to x in Q. Then a is conjugate to x in $Q(n)$ for some n. If $n = 0$, then $a = x \in N$ since $Q(0) = A$ is abelian. If $n > 0$, without loss of generality we may assume that a is not conjugate to x in $Q(n-1)$. Then $a = x_{f(n)}$ by the Collins Lemma [56, Theorem 2.5, Ch. IV]. □

Finally we embed Q into a finitely presented group H such that the word problem is decidable in H and two elements of Q are conjugate in H if and only if they are conjugate in Q. (Such an embedding exists by the Olshanskii–Sapir Theorem, see [66].)

It follows from the construction that an element x_i is conjugate to x in H if and only i belongs to the range of f. As the range of f is not recursive, the problem of whether a given element $h \in H$ is conjugate to x in H is unsolvable. □

5.3 Algorithmic problems for hyperbolic elements

In this section we assume that G is a finitely generated group hyperbolic relative to a collection $\{H_1, \ldots, H_m\}$ of subgroups. We also use some notation introduced at the beginning of Chapter 3. For two elements $f, g \in G$ we write $f \sim g$ if f is conjugate to g.

Recall that the conjugacy problem for a group G given by a recursive presentation is to decide, for any two elements g_1 and g_2 of G, whether or not g_1 is conjugate to g_2 in G. Recently Bumagin [20] proved the following.

Theorem 5.12 (Bumagin, [20]). *Suppose that a finitely generated group G is hyperbolic relative to recursively presented subgroups $\{H_1, \ldots, H_m\}$ and the conjugacy problem is solvable for each of the subgroups $H_1, \ldots, H_m$. Then the conjugacy problem is solvable for G.*

By the *conjugacy problem for hyperbolic elements* we mean the following: given two hyperbolic elements $f, g \in G$, decide whether $f \sim g$. Observe that the next theorem is true without any assumptions about the conjugacy problem in $H_1, \ldots, H_m$.

Theorem 5.13. *Suppose that the word problem is solvable for all H_i, $i = 1, \ldots, m$. Then the conjugacy problem for hyperbolic elements is solvable in G.*

Proof. Since the word problem is solvable in G by Corollary 5.5 , it is enough to show that there is a recursive function $\alpha : \mathbb{N} \to \mathbb{N}$ such that for any two conjugate hyperbolic elements $f, g \in G$ satisfying the inequality $\max\{|f|_X,\ |g|_X\} \leq k$, there exists an element $t \in G$ such that $|t|_X \leq \alpha(k)$ and $f^t = g$.

Let (p, q) be a minimal pair of symmetric geodesics in $\Gamma(G, X \cup \mathcal{H})$ with characteristic elements f, g. Note that no synchronous components of p and q are connected. Indeed otherwise f and g are parabolic. The rest of the proof almost coincide with the proof of Lemma 5.7. By Lemma 3.39 there are no connected components of p and q at all. Further by Corollary 3.42 the length of the element $t = \overline{\phi(p)} = \overline{\phi(q)}$ is not greater than $\rho(k)$. Note that p and q are k–connected geodesics. By Theorem 3.23, any component s of p has X–length at most $|s|_X \leq \varepsilon(1, 0, k)$. Thus we have $|t|_X \leq \alpha(k)$ for $\alpha(k) = \rho(k)\varepsilon(1, 0, k)$. $\square$

Our results concerning algebraic properties of relatively hyperbolic groups allow to treat 'relative versions' of some other algorithmic problems.

Definition 5.14. The *order problem for hyperbolic elements* is to calculate the order of a given hyperbolic element. The *root problem for hyperbolic elements* is to decide whether for a given hyperbolic element $g \in G$ there exists $f \in G$ and $n \in \mathbb{N}$, $n > 1$, such that $g = f^n$. The *power conjugacy problem for hyperbolic elements* is to decide whether for given hyperbolic elements $f, g \in G$ there exist $k, l \in \mathbb{Z}$, such that g^k and f^l are hyperbolic and $g^k \sim f^l$.

For known results about the ordinary order, power conjugacy, and root problems in various classes of groups and relations between these problems we refer to [7, 24, 26, 38, 60, 53, 54, 55, 57].

Theorem 5.15. *Suppose that the word problem is solvable in H_i for any $i = 1, \ldots, m$. Then the order problem for hyperbolic elements is solvable in G.*

Proof. Given a hyperbolic element $g \in G$, there are only finitely many possibilities for the order of g by Theorem 4.2. Thus the order problem in G is reduced to the word problem. $\square$

The next lemma will help us to treat the root problem.

Lemma 5.16. *There exists a recursive function $\beta : \mathbb{N} \to \mathbb{N}$ satisfying the following condition. Let g be an element of G, $|g|_X = k$, and let f be a hyperbolic element of g such that $f^n = g$ for some $n \in \mathbb{N}$. Then f is conjugate to an element $f_0 \in G$ such that $|f_0|_X \leq \beta(k)$.*

Proof. Let f_0 be the element with minimal relative length in the conjugacy class f^G, U a shortest word in $X \cup \mathcal{H}$ representing f_0. By t we denote an element of G such that $f_0 = f^t$. Let T, S be the shortest words in X representing t and g respectively. For any $j \in \mathbb{N}$, there is a cycle $c_j = p_j q_j^{-1}$ in $\Gamma(G, X \cup \mathcal{H})$ such that $\phi(p_j) \equiv U^{nj}$, $\phi(q_j) \equiv T^{-1} S^j T$, as $f_0^{nj} = (g^j)^t$.

If $|f_0|_{X\cup\mathcal{H}} \geq 8\delta + 1$, where δ is the hyperbolicity constant of $\Gamma(G, X \cup \mathcal{H})$, then p_j is $(3, 2\delta)$–quasi–geodesic by Lemma 4.3. We obtain

$$|t^{-1}g^j t|_{X\cup\mathcal{H}} = dist_{X\cup\mathcal{H}}((p_j)_-, (p_j)_+) \geq \frac{1}{3}(l(p_j) - 2\delta) \geq \frac{j}{3}|f_0|_{X\cup\mathcal{H}} - 2\delta.$$

Thus

$$j|g|_{X\cup\mathcal{H}} + 2|t|_{X\cup\mathcal{H}} \geq \frac{j}{3}|f_0|_{X\cup\mathcal{H}} - 2\delta.$$

Dividing by j and passing to the limit as $j \to \infty$, we obtain

$$|f_0|_{X\cup\mathcal{H}} \leq 3|g|_{X\cup\mathcal{H}} \leq 2|g|_X \leq 3k.$$

Thus in any case we have

$$|f_0|_{X\cup\mathcal{H}} \leq \max\{8\delta + 1,\, 3k\}. \tag{5.4}$$

Suppose that for some $j \in \mathbb{N}$, there exist two connected components of p_j, i.e., $p_j = as_1p_0s_2c$, where s_1, s_2 are connected components of p_j. Without loss of generality we may assume that no components of p_0 are connected. Note that f is hyperbolic. Repeating the arguments from the proof of Lemma 4.5 and using (5.4), we obtain the estimate

$$|f_0|_X \leq 2ML(\max\{8\delta + 1,\, 3k\})^2. \tag{5.5}$$

Further assume that for any $j \in \mathbb{N}$, all components of p_j are isolated in p_j. This yields that all components of c_j are isolated in c_j since $\phi(q_j)$ is a word in the alphabet X. By Lemma 3.1, for any $i = 1, \ldots, m$, any H_i–syllable V of U satisfies the inequality

$$j|n||\overline{V}|_X \leq MLl(c_j) \leq ML(j|n||f_0|_{X\cup\mathcal{H}} + 2|t|_X + j|g|_X).$$

Thus

$$|\overline{V}|_X \leq ML\left(|f_0|_{X\cup\mathcal{H}} + \frac{2}{j}|t|_X + k\right).$$

Assuming $j \to \infty$, we obtain

$$|\overline{V}|_X \leq ML\left(|f_0|_{X\cup\mathcal{H}} + k\right) \leq 2ML\max\{8\delta + 1,\, 3k\}.$$

This implies (5.5) again. Thus we can set

$$\beta(k) = 2ML(\max\{8\delta + 1,\, 3k\})^2.$$

□

Theorem 5.17. *Suppose that the word problem is solvable in H_i for any $i = 1, \ldots, m$. Then the root problem for hyperbolic elements is solvable in G.*

Proof. Let g be a hyperbolic element of G. Since the order problem for hyperbolic elements is solvable in G, we can decide whether the order of g is finite. Let us consider two cases.

1. The order of g is finite. Recall that the set of powers of hyperbolic elements of G is finite. Let n_0 be the maximal finite order of hyperbolic elements in G. If $f^n = g$ for some positive n, then we may assume that $n \leq n_0$. By Lemma 5.16, to decide whether g has a non–trivial root in G, it suffices to decide whether there exists an element $f_0 \in G$ and a natural number $n \leq n_0$ such that $|f_0|_X \leq \beta(|g|_X)$ and $f_0^n \sim g$. Thus the root problem is reduced to the conjugacy problem for hyperbolic elements, which is solvable by Theorem 5.13.

2. The order of g is infinite. If $f^n = g$ for some $f \in G$, $n \in \mathbb{N}$, then

$$\tau^{rel}(f) = \frac{1}{n}\tau(g) \leq \frac{1}{n}|g|_{X\cup\mathcal{H}} \leq \frac{1}{n}|g|_X.$$

Thus, by Theorem 4.25, $n \leq |g|_X/d$, where d is a constant which is independent of g and f. Using Lemma 5.16, we now can reduce the root problem to the conjugacy problem for hyperbolic elements as in the first case. □

To deal with the power conjugacy problem, we need the following.

Lemma 5.18. *There exists a recursive function* $\gamma : \mathbb{N} \to \mathbb{N}$ *such that if* $f, g \in G$ *are two hyperbolic elements of infinite order and* $f^k \sim g^l$ *in* G *for some* $k, l \in \mathbb{Z} \setminus \{0\}$*, then there exist* $k', l' \in \mathbb{Z} \setminus \{0\}$ *such that* $f^{k'} \sim g^{l'}$ *in* G *and*

$$\max\{|k'|,\ |l'|\} \leq \gamma(\max\{|f|_X,\ |g|_X\}).$$

Proof. Suppose that k and l are numbers with minimal $\max\{|k|,\ |l|\}$ among all pairs k, l satisfying $f^k \sim g^l$. We are going to show that

$$\max\{|k|,\ |l|\} \leq \gamma(\max\{|f|_X,\ |g|_X\}) \tag{5.6}$$

for some recursive function γ. For definiteness we assume that $|k| \geq |l|$.

Let t be an element of G such that

1) $g^l = tf^kt^{-1}$ and

2) t has minimal relative length among all elements of G satisfying 1).

Let U, V (respectively T) be shortest words over X (respectively $X \cup \mathcal{H}$) representing elements f, g (respectively t). The conditions of the lemma imply that there exists a quadrangle

$$rps^{-1}q^{-1}$$

in $\Gamma(G, X \cup \mathcal{H})$ such that $\phi(r) \equiv \phi(s) \equiv T$, $\phi(p) \equiv U^k$, $\phi(q) \equiv V^l$. For convenience, we assume $r_- = 1$.

Let $a_1, \ldots a_k$ denote the ending vertices of the subpaths $p_1, \ldots, p_k$ of p with $(p_i)_- = p_-$ and $\phi(p_i) \equiv U^i$, $i = 1, \ldots, k$. Let $\lambda > 0$ be a number such that there is no hyperbolic element h of infinite order in G with $\tau^{rel}(h) < \lambda$. Note that p and q are (λ, c)–quasi–geodesics in $\Gamma(G, X \cup \mathcal{H})$, for $c = \max\{|f|_X,\ |g|_X\}$ since



Figure 5.1:

$\tau^{rel}(g) = \inf\{|g^n|_{X\cup\mathcal{H}}/n\}$. By Corollary 3.33 and Theorem 3.23, for any vertex $a_i \in p$ there exists a vertex $b_i \in q \cup r \cup s$ such that

$$dist_X(a_i, b_i) \le 2(\nu + \varepsilon), \tag{5.7}$$

where $\varepsilon = \varepsilon(\lambda, c, 0)$. First of all we wish to show that if a_i is sufficiently far from the endpoints of p, then $b_i \in q$.

For instance, suppose that $b_i \in r$. Let o be a geodesic in $\Gamma(G, X \cup \mathcal{H})$ such that $o_- = b_i$, $o_+ = a_i$ (see Fig. 5.1). From (5.7), we obtain $l(o) \le 2(\nu + \varepsilon)$. Note that the element a_i satisfies $g^l a_i f^k a_i^{-1}$ since $a_i = tf^i$. By the choice of t this means that

$$|a_i|_{X\cup\mathcal{H}} \ge |t|_{X\cup\mathcal{H}}. \tag{5.8}$$

Notice that

$$|a_i|_{X\cup\mathcal{H}} \le |b_i|_{X\cup\mathcal{H}} + l(o) \le |b_i|_{X\cup\mathcal{H}} + 2(\nu + \varepsilon) \tag{5.9}$$

and

$$|t|_{X\cup\mathcal{H}} = |b_i|_{X\cup\mathcal{H}} + l([b_i, t]) \tag{5.10}$$

as r is geodesic. Combining (5.8), (5.9), and (5.10), we obtain the following estimate on the length of the segment $[b_i, t]$ of r:

$$l([b_i, t]) \le 2(\nu + \varepsilon).$$

Hence,

$$dist_{X\cup\mathcal{H}}(t, a_i) \le dist_{X\cup\mathcal{H}}(t, b_i) + dist_{X\cup\mathcal{H}}(b_i, a_i) \le 4(\nu + \varepsilon).$$

Since p is (λ, c)–quasi–geodesic, we have

$$l(p_i) \le \lambda dist_{X\cup\mathcal{H}}(t, a_i) + c \le 4\lambda(\nu + \varepsilon) + c.$$

Therefore, if $i \ge N$ for

$$N = 4\lambda(\nu + \varepsilon) + c,$$

then b_i can not belong to r as $l(p_i) = \|U\| i \ge i$. Similarly, if $i < k - N$, then b_i can not belong to s.

Suppose that $k > 2N$. Then for any $N \leq i \leq k - N$, $b_i \in q$. We denote by c_j the terminal vertex of the subpath q_j of q, such that $(q_j)_- = q_- = 1$ and $\phi(q_j) \equiv V^j$. Thus $c_j = g^j$. Let $c_{j(i)}$ be the closest vertex (with respect to the relative metric) to the vertex b_i. (Here the index $j(i)$ depends of i.) Obviously

$$dist_X(c_{j(i)}, b_i) \leq \frac{1}{2}|g|_X.$$

Hence,

$$|a_i^{-1}c_{j(i)}|_X = dist_X(a_i, c_{j(i)}) \leq \frac{1}{2}|g|_X + 2(\nu + \varepsilon).$$

Let

$$k_0 = 2N + (card\ X)^{\frac{1}{2}|g|_X + 2(\nu+\varepsilon)} + 1.$$

If $k > k_0$, then there exists two pairs $a_{i_1}, c_{j(i_1)}$ and $a_{i_2}, c_{j(i_2)}$ such that $a_{i_1}^{-1}c_{j(i_1)} = a_{i_2}^{-1}c_{j(i_2)}$ since the number of different elements of G of X–length at most n is less that or equal to $(card\, X)^n + 1$. Let $t_0 = a_{i_1}^{-1}c_{j(i_1)}$, $j_0 = j(i_1) - j(i_2)$, and $i_0 = i_1 - i_2$. Then t_0^{-1} conjugate g^{j_0} to f^{i_0} or f^{-i_0}. Since $|i_0| \leq |k|$, we arrive to the contradiction with the choice of k and l. Thus k is less than or equal to k_0 and the inequality (5.6) is true for

$$\gamma = 2N + (card\ X)^{\frac{1}{2}|g|_X + 2(\nu+\varepsilon)} + 1.$$

□

Theorem 5.19. *Suppose that the word problem is solvable in H_i for any $i = 1, \ldots, m$. Then the power conjugacy problem for hyperbolic elements is solvable in G.*

Proof. Let f, g be two hyperbolic elements of G. There are three cases to consider (by Theorem 5.15 they can be effectively recognized).

1. Both the elements f, g have finite orders n_1 and n_2 respectively. Then it suffices to decide whether the elements f^k and g^l are hyperbolic and conjugate for some k and l satisfying $0 < k < n_1$, $0 < l < n_2$. This can be done by Theorems 5.6 and 5.13.

2. g has finite order, f has infinite order (or conversely). Clearly f is not conjugate to g in this case.

3. Both the elements f, g have infinite order. Then applying Lemma 5.18, we can reduce the question to the conjugacy problem for hyperbolic elements in G as in the first case. □

Open questions

Here we state some natural problems and conjectures which seem to be important for the further studying relatively hyperbolic groups.

If G is a finitely presented group and is hyperbolic relative to a subgroup H, then H is finitely generated by Proposition 2.29. However the following important question remains open.

Problem 5.1. *Let G be a finitely presented group hyperbolic relative to a subgroup H. Does it follow that H is finitely presented?*

Assume that G is a group generated by a finite set X hyperbolic relative to a collection of subgroups $H_1, \ldots, H_m$. As we have already mentioned in Section 4.2, Theorem 4.16 does not hold without the assumption *card* $(R \cap H_i^g) < \infty$. However, one can try to prove a similar result in the general case. More precisely, let R be a relatively quasi–convex subgroup of G. We consider the set of subgroups

$$\mathfrak{S} = \{H_i^g \cap R \mid i = 1, \ldots, m,\ g \in G\}.$$

The subgroup R acts on $\mathfrak{S}$ by conjugations. If R is quasi–convex, it is not hard to show that the number of orbits of this action is finite. Let $P_1, \ldots, P_l$ be representatives of the orbits.

Problem 5.2. *Prove that the group R is relatively hyperbolic with respect to $\{P_1, \ldots, P_l\}$.*

Note that R and P_i need not be finitely generated in this case. The simplest example of this type is the pair $G = H_1 * H_2$ and $R = \langle K_1, K_2 \rangle$ considered in Remark 4.17. The proof of the conjecture should be slightly more complicated than one of Theorem 4.16, although the difficulties are rather technical.

Another problem about relatively quasi–convex groups is

Problem 5.3. *Does the notion of relative quasi–convexity formulated in this paper coincide with the dynamical quasi–convexity introduced by Bowditch in [**?**]?*

Let G be a group generated by a finite set X, $H_1, \ldots, H_m$ subgroups of G. We consider the following condition:

($*$) *There exists a constant $\nu > 0$ such that for any geodesic triangle $\Delta = pqr$ in $\Gamma(G, X \cup \mathcal{H})$ and any vertex v on p, there exists a vertex u on the union $q \cup r$ such that $dist_X(u, v) \leq \nu$.*

Theorem 3.26 shows that if G is hyperbolic relative to $\{H_1, \ldots, H_m\}$, then ($*$) is satisfied. The converse is not true. Indeed any group G satisfies ($*$) with respect to any subgroup H of finite index. However, by Proposition 2.36, G is never hyperbolic relative to H unless $G = H$ or H is finite. On the other hand, it is easy to see that any group G satisfying ($*$) is weakly hyperbolic (or hyperbolic in the sense of Farb) relative to $H_1, \ldots, H_m$. And again the converse is not true. For example, if $G = H \times \mathbb{Z}$, where H is an infinite finitely generated group, then G is weakly hyperbolic relative to H, but do not satisfy ($*$). Thus the class of groups satisfying ($*$) is intermediate between the classes of hyperbolic and weakly hyperbolic groups.

Problem 5.4. *Study the class of finitely generated groups satisfying ($*$) with respect to a finite collection of subgroups.*

The next two problems are inspirited by well–known questions about ordinary hyperbolic groups.

Problem 5.5. *Assume that G is a finitely generated group hyperbolic relative to a collection of subgroups $\{H_1, \ldots, H_m\}$. Suppose that all subgroups H_i are Hopfian. Does it follow that G is Hopfian?*

Recall that a group G is said to be Hopfian if every epimorphism $G \to G$ is an isomorphism. First examples of non–Hopfian groups were found by Boumslag and Solitar [5]: the group

$$BS(m, n) = \langle a, t \mid (a^m)^t = a^n \rangle, \tag{5.11}$$

is not Hopfian whenever m and n are relatively prime and neither of $|m|$, $|n|$ is equal to 1. We note that $BS(m, n)$ is weakly relatively hyperbolic with respect to $\langle a \rangle$ (see [67]). Sela proved that if a torsion–free hyperbolic group does not decompose as a free product, then it is Hopfian. It seems to be reasonable to assume that analogous result is true in the relative case.

Problem 5.6. *Assume that G is a group hyperbolic relative to a collection of subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$. Suppose that all subgroups H_λ are residually finite. Can G be non–residually finite?*

This question is open even for ordinary hyperbolic group G. It is known [45, 63] that an infinite hyperbolic group is never simple, but it is still unknown whenever it is always residually finite (some speculations can also be found in [49], [64], and [51]). One of the results in this direction can be found in the paper [83], where Wise showed that negatively curved polygons of finite groups are residually finite. We note that in the case of CAT(0)–groups the answer to Problem 5.6 is known to be positive. In the paper [82], Wise produced examples of compact non–positively curved spaces whose fundamental groups are not residually finite. Subsequently, Burger and Mozes [22] constructed compact non–positively curved 2–complexes whose fundamental groups are simple.

Appendix. Equivalent definitions of relative hyperbolicity

The definition of Bowditch

The original definition of Bowditch characterizes relative hyperbolicity in dynamical terms.

Definition 6.1. A finitely generated group G is *hyperbolic relative to a collection of finitely generated subgroups* $H_1, \ldots, H_m$ if it admits a properly discontinuous isometric action on a path–metric hyperbolic proper space X such that the induced action of G on the boundary ∂X satisfies the following conditions.

(1) G acts on ∂X as a geometrically finite convergence group.

(2) The maximal parabolic subgroups of G are precisely the subgroups of G conjugate to $H_1, \ldots, H_m$.

Convergence groups were introduced by Gehring and Martin [40] in order to describe the dynamical properties of Kleinian groups acting on the standard sphere in $\mathbb{R}^n$ and were generalized to groups acting on compact Hausdorff spaces by Tukia and Freden [81], [39]. Their motivation came from the observation that an isometry group of a hyperbolic space X acts as a convergence group on the hyperbolic boundary of X.

We recall that a group G of homeomorphisms of a metrizable compactum M acts on M as a *convergence group* if the induced action on the space of distinct triples of elements of M is properly discontinuous (for equivalent definitions we refer to [**?**]). One says that a subgroup $H \leq G$ is *parabolic* if it is infinite, fixes some point of M and contains no elements $g \in G$ of infinite order such that $card(fix(g)) = 2$. In this case the fixed point of H is unique and is called a *parabolic point.* A parabolic point x is said to be *bounded* if $(M \setminus \{x\}) / Stab_G(x)$ is compact. A point $x \in M$ is called a *conical limit point* if there is a sequence $\{g_i\}$ and two distinct points $a, b \in M$ such that $g_i x$ converges to a and $g_i y$ converges to b for any $y \in M \setminus \{x\}$. Finally, a convergence group G is said to be *geometrically finite* if every point of M is a conical limit point or a bounded parabolic point.

In [13], Bowditch proved that Definition 6.1 is equivalent to the following.

Definition 6.2. A finitely generated group G is *hyperbolic relative to a collection of finitely generated subgroups* $H_1, \dots, H_m$ if it admits an action on a hyperbolic graph K such that the following condition hold.

1) All edge stabilizers are finite.

2) All vertex stabilizers are finite or conjugate to one of the subgroups $H_1, \dots, H_m$.

3) The number of orbits of edges is finite.

4) The graph K is fine, that is, for every $n \in \mathbb{N}$, any edge of K is contained in finitely many circuits of length n. (Here circuit means a cycle without self–intersections).

The definition of Farb

Let G be a group generated by a finite set X and let $\{H_1, H_2, \dots H_m\}$ be a collection of subgroups of G. We begin with the Cayley graph $\Gamma(G, X)$ of G and form a new graph as follows: for each left coset gH_i of H_i in G, add a vertex $v(gH_i)$ to $\Gamma(G, X)$, and add an edge $e(gh)$ of length $1/2$ from each element gh of gH_i to the vertex $v(gH_i)$. The new graph is called the *coned–off Cayley graph* of G with respect to $\{H_1, H_2, \dots H_m\}$, and is denoted by $\widehat{\Gamma}(G, X)$. We give this graph the path metric. Note that $\widehat{\Gamma}(G, X)$ is not a proper metric space as closed balls are not necessarily compact.

Definition 6.3. The group G is *hyperbolic relative to* $\{H_1, H_2, \dots H_m\}$ if the coned–off Cayley graph $\widehat{\Gamma}(G, X)$ of G with respect to $\{H_1, H_2, \dots H_m\}$ is a hyperbolic metric space.

Definition 6.4. Given a path p in $\widehat{\Gamma}(G, X)$, we say that p *penetrates* the coset gH_i if p passes through the cone point $v(gH_i)$; a vertex v_1 (respectively v_2) of the path p which precedes to $v(gH_i)$ (respectively succeeds to $v(gH_i)$) is called an *entering vertex* (respectively an *exiting vertex*) of p in the coset gH_i. Notice that entering and exiting vertices are always vertices of $\Gamma(X, G)$. A path p in $\widehat{\Gamma}(G, X)$ is said to be a path *without backtracking* if, for every coset gH_i which p penetrates, p never returns to gH_i after leaving gH_i.

Definition 6.5. (Bounded coset penetration). The pair $(G, \{H_1, H_2, \dots H_m\})$ is said to satisfy the *Bounded Coset Penetration property* (or BCP property for brevity) if, for every $\lambda \geq 1$, there is a constant $a = a(\lambda) > 0$ such that if p and q are $(\lambda, 0)$–quasi–geodesics without backtracking in $\widehat{\Gamma}(G, X)$ such that the endpoints of p and q are in $\Gamma(G, X)$, $p_- = q_-$, and $dist_X(p_+, q_+) \leq 1$, then the following conditions hold.

(1) If p penetrates a coset gH_i but q does not penetrate gH_i, then the

entering vertex and the ending vertex of p in gH_i are an X–distance of at most a from each other.

(2) If both p and q penetrate a coset gH_i, then the entering vertices of p and q in gH_i lies an X–distance of at most a from each other; similarly for the exiting vertices.

Example 6.6. The group $G = \langle a, b \mid [a, b] = 1\rangle \cong \mathbb{Z} \times \mathbb{Z}$ is weakly hyperbolic relative to the cyclic subgroup $H = \langle a\rangle$. However the pair (G, H) does not have the BCP property: the paths a^n and ba^n are relative geodesics without backtracking ending a distance 1 apart in Γ, but they clearly violate condition (1) of Definition 6.5 when n is large enough.

Dahmani [31] shows that G satisfies Definition 6.2 if and only if it satisfies Definition 6.3 and the pair $(G, \{H_1, H_2, \ldots H_m\})$ has the BCP property. (The proof of this fact in [30] contains gaps and wrong statements.) The proof can also be found in [21]. Now we are going to reformulate Farb's definition in terms of the relative Cayley graph $\Gamma(G, X \cup \mathcal{H})$.

Definition 6.7. Two metric spaces M_1, M_2 are said to be *quasi–isometric* if there exist $\lambda > 0$, $c \geq 0$, $\varepsilon \geq 0$, and a map $\alpha : M_1 \to M_2$ such that the following two condition hold.

1. For any $x, y \in M_1$, we have
$$\frac{1}{\lambda} dist_{M_1}(x, y) - c \leq dist_{M_2}(\alpha(x), \alpha(y)) \leq \lambda dist_{M_1}(x, y) + c.$$
2. For any $z \in M_2$ there exists $x \in M_1$ such that
$$dist_{M_2}(\alpha(x), z) \leq \varepsilon.$$

Recall that hyperbolicity of metric spaces is invariant under quasi–isometry.

Lemma 6.8. *Let G be a group, $\{H_1, H_2, \ldots, H_m\}$ a finite collection of subgroups of G. The conned–off Cayley graph $\widehat{\Gamma}$ is quasi–isometric to $\Gamma(G, X \cup \mathcal{H})$ endowed with the relative metric $dist_{X\cup\mathcal{H}}$. In particular, $\widehat{\Gamma}$ is hyperbolic if and only if so is $\Gamma(G, X \cup \mathcal{H})$.*

Proof. Note that the identity map on G induces in isometric embedding ι of the vertex set $V(\Gamma(G, X \cup \mathcal{H}))$ of $\Gamma(G, X \cup \mathcal{H})$ to $\widehat{\Gamma}$ and $\widehat{\Gamma}$ belongs to the closed 1–neighborhood of the image $\iota(V(\Gamma(G, X \cup \mathcal{H})))$. □

Clearly, the BCP property can be rewritten as follows (see Section 2.2 for necessary definitions).

Lemma 6.9. *Let G be a group generated by a finite set X, $\{H_1, H_2, \ldots, H_m\}$ a finite collection of finitely generated subgroups of G. The pair $(G, \{H_1, H_2, \ldots, H_m\})$ satisfies the BCP property if and only if for any $\lambda \geq 1$,* *there exists constant $a = a(\lambda)$ such that the following conditions hold. Let p, q be $(\lambda, 0)$–quasi–geodesics without backtracking in $\Gamma(G, X \cup \mathcal{H})$ (in the sense of Definition 3.9) such that $p_- = q_-$, $dist_X(p_+, q_+) \le 1$.*

1) Suppose that for some i, s is an H_i–component of p such that $dist_X(s_-, s_+) \ge a$; then there exists an H_i–component t of q such that t is connected to s.

2) Suppose that for some i, s and t are connected H_i–components of p and q respectively. Then $dist_X(s_-, t_-) \le a$ and $dist_X(s_+, t_+) \le a$.

Proof of the main theorem

Our main goal here is to prove

Theorem 6.10. *Let G be a group generated by a finite set X, $\{H_1, H_2, \ldots, H_m\}$ a collection of subgroups of G. Then the following conditions are equivalent.*

1) G is relatively finitely presented with respect to $\{H_1, H_2, \ldots, H_m\}$ and the relative Dehn function of the pair $(G, \{H_1, H_2, \ldots, H_m\})$ is linear.

2) G is hyperbolic with respect to the collection $\{H_1, H_2, \ldots, H_m\}$ in the sense of Farb and satisfies the BCP property (or, equivalently, G is hyperbolic with respect to $\{H_1, H_2, \ldots, H_m\}$ in the sense of Bowditch).

Proof. Theorem 3.23 gives the implication 1) ⇒ 2). Let us show that Farb's definition implies relative hyperbolicity in the sense of our paper.

Let p be a cycle in $\Gamma(G, X \cup \mathcal{H})$. We say that p is *atomic* if any subpath q of p of length $l(q) \le 1/2l(p)$ is geodesic in $\Gamma(G, X \cup \mathcal{H})$. Recall that $\Gamma(G, X \cup \mathcal{H})$ is hyperbolic by Lemma 6.8. In what follows we denote by δ the hyperbolicity constant of $\Gamma(G, X \cup \mathcal{H})$.

Lemma 6.11. *Let p be an atomic cycle in $\Gamma(G, X \cup \mathcal{H})$. Then the following conditions hold.*

1) $l(p) \le 4\delta + 9$.

2) For any $i = 1, \ldots, m$, any H_i–component of p is isolated.

Proof. 1) Suppose that $l(p) > 4\delta + 9$. We can represent p as the product $p = p_1p_2p_3$, where $l(p_1) = l(p_2)$ and $l(p_3) \le 1$. Then $p_1p_2p_3$ is a geodesic triangle in $\Gamma(G, X \cup \mathcal{H})$ since p is atomic. At least one of the sides p_1, p_2 is longer than $2\delta + 4$. Assume that $l(p_1) > 2\delta + 4$. Let us take the middle point w of p_1. Since $\Gamma(G, X \cup \mathcal{H})$ is δ–hyperbolic, there exists a point $z \in p_2p_3$ such that $dist_{X \cup \mathcal{H}}(w, z) \le \delta$. Let v (respectively u) be a vertex of p_1 (respectively p_2p_3) that is closest to w (respectively z). Then $dist_{X \cup \mathcal{H}}(v, u) \le \delta + 1$. Notice that both the segments $[u, v]$ and $[v, u]$ of p are not geodesic as both of them have length at least $l(p_1)/2 - 1/2 > \delta + 3/2$. However, at least one of these segments has length at most $1/2l(p)$ contradictory to the assumption that p is atomic.

2) Let $p = arbs$, where a, b are connected H_i–components of p. Then both the subpaths ar and bs are not geodesic since

$$dist_{X\cup\mathcal{H}}(a_-, r_+) = dist_{X\cup\mathcal{H}}(b_-, s_+) = 1.$$

Thus we get a contradiction as above. $\square$

Corollary 6.12. *Let $\mathcal{A}$ be the set of labels of all atomic cycles in $\Gamma(G, X\cup\mathcal{H})$. Then card $\mathcal{A} < \infty$.*

Proof. Let p be an atomic cycle. We can represent p as $p = cd$, where $l(c) \le l(d) \le l(c) + 1$. Since p is atomic, c is geodesic and d is $(2, 0)$–quasi–geodesic as any proper subpath of d is geodesic. Since all components of p are isolated, c and d are paths without backtracking. Therefore, for any i, the X–length of every H_i–component of p is at most a, where $a = a(2)$ is the constant provided by Lemma 6.9. Hence there are only finitely many possibilities for labels of H_i–components of atomic cycles. This fact together with the first assertion of Lemma 6.11 implies the finiteness of $\mathcal{A}$. $\square$

Let us return to the proof of the theorem. We are going to show that G has the relative presentation

$$\langle X, \mathcal{H} \mid R = 1,\ R \in \mathcal{A}\rangle$$

and the corresponding relative Dehn function satisfies $\delta^{rel}(n) \le 2^n$. Let W be a word of length at most n over $(X\cup\mathcal{H})^*$ representing 1 in G. We have to show that there exists a van Kampen diagram with boundary label W having at most 2^n cells labelled by words from $\mathcal{A}$.

We proceed by induction on n. Consider the cycle p in $\Gamma(G, X\cup\mathcal{H})$ labelled W. If p is atomic, the existence of the required diagram is obvious. If p is not atomic, then without loss of generality we can assume that p is combinatorially homotopic to a product q_1q_2 of cycles q_1, q_2 of length $l(q_i) \le l(p) - 1$, $i = 1, 2$. This means that up to a cyclic shift $W =_F V_1V_2$, where $\|V_i\| \le \|W\| - 1 \le n - 1$, and V_i represents 1 in G for $i = 1, 2$. By the inductive assumption, there exist van Kampen diagrams with boundary labels V_1, V_2 and with the number of cells labelled by words from $\mathcal{A}$ at most $2^{\|V_1\|}$ and $2^{\|V_1\|}$ respectively. Gluing these diagrams together in the obvious way, we obtain a van Kampen diagram with boundary label W and with the number of cells labelled by words from $\mathcal{A}$ at most

$$2^{\|V_1\|} + 2^{\|V_2\|} \le 2^{n-1} + 2^{n-1} = 2^n.$$

Thus we proved that the relative Dehn function of G with respect to $\{H_1, \ldots, H_m\}$ is well–defined. Since $\Gamma(G, X\cup\mathcal{H})$ is hyperbolic, the relative Dehn function is, in fact, linear by Corollary 2.54. $\square$

Bibliography

[1] A.D. Alexandrov, *Rulled surfaces in metric spaces*, Vestnik Leningrad. Univ., **12** (1957), 5–26.

[2] E. Alibegovic, *A Combination Theorem for Relatively Hyperbolic Groups Authors,* prep., 2003, available at http://xxx.lanl.gov/abs/math.GR/0310257.

[3] J.M. Alonso, M.R. Bridson, *Semyhyperbolic groups*, Proc. London Math. Soc., (3) **70** (1995), 56–114.

[4] P. Bahls, *Relative hyperbolicity and right-angled Coxeter groups,*, prep., 2004, available at http://xxx.lanl.gov/abs/math.GR/0401280.

[5] G. Baumslag, D. Solitar, *Some two-generator one-relator non-Hopfian groups*, Bull. Amer. Math. Soc. **68** (1962) 199–201.

[6] V.N. Berestovskii, I.G. Nikolaev, *Multidimensional generalized Riemannian spaces,* Geometry IV (Yu. Reshetnyak, ed.), Encyclopedia of Math. Sci., Vol. 70, Springer–Verlag, 1993, 165–243.

[7] V.N. Bezverkhnii, V.A. Grinblat, *The root problem in Artin groups* (Russian), Algorithmic problems of the theory of groups and semigroups, pp. 72–81, Tulsk. Gos. Ped. Inst., Tula, 1981

[8] O.V. Bogopolskii, V.N. Gerasimov, *Finite subgroups of hyperbolic groups,* Algebra and Logic **34** (1995), 343–345.

[9] B.H. Bowditch, *Notes on Gromov's hyperbolicity criterion for path–metric spaces,* Group Theory from a Geometrical Viewpoint (E. Ghys, A. Haefliger, A. Verjovsky, ed.), Proc. ICTP Trieste 1990, Word Scientific, Singapore, 1991, 373–464.

[10] B.H. Bowditch, *A short proof that a subquadratic isoperimetric inequality implies a linear one,* Michigan J. Math., **42** (1995), 103–107.

[11] B.H. Bowditch, *A topological characterization of hyperbolic groups,* J. Amer. Math. Soc., **11** (1998), 643–667.

[12] B.H. Bowditch, *Convergence groups and configuration spaces,* In: Group Theory Down Under (ed. J.Cossey, C.F. Miller, W.D. Neumann, M. Shapiro), de Gruyter, 1999, 23–54.

[13] B.H. Bowditch, *Relatively hyperbolic groups,* prep., 1999.

[14] B.H. Bowditch, *Intersection numbers and the hyperbolicity of the complex of curves,* prep., 2002.

[15] S.G. Brick, J.M. Corson, *Annular Dehn functions of groups,* Bull. Austral. Math. Soc. 58 (1998), no. 3, 453–464.

[16] S.G. Brick, J.M. Corson, *Dehn functions and complexes of groups,* Glasgow Math. J., **40** (1998), 33–46.

[17] S.G. Brick, J.M. Corson, *On Dehn functions of amalgamations and strongly undistorted subgroups,* IJAC, **10** (2000), 5, 665–681.

[18] M. Bridson, *Polynomial Dehn functions and the length of asynchronously automatic structures,* Proc. LMS, to appear.

[19] M. Bridson, A. Haefliger, Metric spaces of non–positive curvature, Springer, 1999.

[20] I. Bumagin, *Conjugacy problem for relatively hyperbolic groups*, submitted to Alg. Geom. Topology.

[21] I. Bumagin, *On the definition of relatyvely hyperbolic groups,* prep., 2004, available at http://xxx.lanl.gov/abs/math.GR/0402072.

[22] M. Burger, S. Mozes, *finitely presented simple groups and products of trees*, C. R. Acad. Sci. Paris, **324** (1997), 747–752.

[23] R. Camm, *Simple free products,* J. London Math. Soc., **28** (1953), 66-76.

[24] F.B. Cannonito, R.W. Gatterdam, *The word problem and power problem in 1-relator groups are primitive recursive*, Pacific J. Math. **61** (1975), no. 2, 351–359.

[25] D.J. Collins, *The word, power and order problems in finitely presented groups,* Word problems: decision problems and the Burnside problem in group theory (Conf., Univ. California, Irvine, Calif., 1969; dedicated to Hanna Neumann), 401–420. Studies in Logic and the Foundations of Math., Vol. 71, North-Holland, Amsterdam, 1973

[26] L.P.J. Comerford, *A note on power-conjugacy,* Houston J. Math. **3** (1977), no. 3, 337–341.

[27] M. Coornaert, T. Delzant, A. Papadopoulos, Notes sur les groupes hyperboliques de Gromov, Springer LNM 1441, 1990.

[28] J.M. Corson, *Groups acting on complexes and complexes of groups,* Geometric Group Theory (Charney, Davis and Shapiro, eds.), Walter de Gruyter, Berlin, New York, 1955, 79–97.

[29] J.M. Corson, *Howie diagrams and complexes of groups,* Comm. Alg., **23** (1995), 14, 5221–5242.

[30] F. Dahmani, *Classifying space and boundary for relatively hyperbolic groups*, Proc. of London Math. Soc., to appear.

[31] F. Dahmani, Les groupes relativement hyperboliques et leurs bords, PhD thesises, 2003.

[32] F. Dahmani, *Combination of convergence groups,* prep., 2002.

[33] W. Dicks, M.J. Dunwoody, Groups acting on graphs, Cambridge Univ. Press, Cambridge, 1989.

[34] P. Eberlein, *Lattices in spaces of nonpositive curvature,* Annals of Math., **111** (1980), 435–476.

[35] D.B.A. Epstein, J. Cannon, D.F. Holt, S. Levy, M.S. Paterson, W.P. Thurston, Word processing in groups, Jones and Bartlett, 1992.

[36] B. Farb, *The extrinsic geometry of subgroups and the generalized word problem*, Proc. London Math. Soc. **68** (1994), 3, 577–593.

[37] B. Farb, *Relatively hyperbolic groups,* GAFA, **8** (1998), 810–840.

[38] B. Fine, *On power conjugacy and* SQ-*universality for Fuchsian and Kleinian groups*, Modular functions in analysis and number theory, 41–54, Lecture Notes Math. Statist., 5, Univ. Pittsburgh, Pittsburgh, PA, 1983. .

[39] E.M. Freden, *Properties of convergence groups and spaces*, Conform. Geom. Dynam., **1** (1997), 13–23.

[40] F.W. Gehring, G.J. Martin, *Discrete quasiconformal groups I*, Proc. London Math. Soc., **55** (1987), 331–358.

[41] S. Gersten, *Reducible diagrams and equations over groups,* Essays in Group Theory (S. Gersten, eds.), MSRI Publ., Springer–Verlag, 1987, 15–74.

[42] S.M. Gersten, *Subgroups of word hyperbolic groups in dimension 2,* J. London Math. Soc., **54** (1996), 261–283.

[43] S.M. Gersten, H.Short, *Rational subgroups of biautomatic groups,* Ann. of Math., (2) **134** (1991), 125–158.

[44] E. Ghys, P. de la Harpe, Eds., Sur les groupes hyperboliques d'aprés Mikhael Gromov, Progress in Math., 83, Birkaüser, 1990.

[45] M. Gromov, *Hyperbolic groups,* Essays in Group Theory, MSRI Series, Vol.8, (S.M. Gersten, ed.), Springer, 1987, 75–263.

[46] M. Gromov, *Asymptotic invariants of infinite groups*, Geometric group theory, Vol. 2 (Sussex, 1991), 1–295, London Math. Soc. Lecture Note Ser., 182, Cambridge Univ. Press, Cambridge, 1993.

[47] V.S. Guba, M.V. Sapir, *On Dehn functions of free products of groups,* Proc. Amer. Math. Soc. **127** (1999), 7, 1885–1891.

[48] W.J. Harvey, *Boundary structure of the modular group*, Riemannian Surfaces and Related Topics: Proc. of the 1978 Stony Brook Conference (I. Kra and B. Maskit, eds.), Ann. of Math. Stud. 97, Princeton, 1981.

[49] S.V. Ivanov, A.Yu. Olshanskii, *Hyperbolic groups and their quotients of bounded exponents*, Trans. Amer. Math. Soc. **348** (1996), no. 6, 2091–2138.

[50] K.A. Kalorkoti, *Decision problems in group theory*, Proc. London Math. Soc. (3) **44** (1982), no. 2, 312–332.

[51] I. Kapovich, D.Wise, *The equivalence of some residual properties of word-hyperbolic groups*, J. Algebra **223** (2000), no. 2, 562–583.

[52] I. Kapovich, *Relative hyperbolicity and Artin groups*, prep., 2002.

[53] L. Larsen, *On the computability of conjugate powers in finitely generated Fuchsian groups*, Acta Math. 139 (1977), no. 3-4, 267–291.

[54] S. Lipschutz, M. Lipschutz, *A note on root decision problems in groups*, Canad. J. Math. **25** (1973), 702–705.

[55] S. Lipschutz, Ch.F. Miller III, *Groups with certain solvable and unsolvable decision problems*, Comm. Pure Appl. Math. **24** (1971), 7–15.

[56] R.C. Lyndon, P.E. Shupp, Combinatorial Group Theory, Springer–Verlag, 1977.

[57] I.G. Lysenok, *On some algoritmic properties of hyperbolic groups,* Math. USSR Izv., **35** (1990), 145–163.

[58] W. Magnus, A. Karras, D. Solitar, Combinatorial group theory, Interscience Publ., 1966.

[59] H. Masur, Y. Minsky, *Geometry of complex of curves I: Hyperbolicity,* Invent. Math. 138 (1999), no. 1, 103–149.

[60] C.F. Miller III, On group Theoretic Decision Problems and their Classification, Annals of Math. Stud., **68**, Princeton Univ. Press, Princeton NJ, 1971.

[61] A.Yu. Ol'shanskii, Geometry of defining relations in groups, Kluwer Academic Publisher, 1991.

[62] A.Yu. Ol'shanskii, *Hyperbolicity of groups with subquadratic isoperimetric inequalities*, Internat. J. Algebra Comput., **1** (1991), 281–289.

[63] A.Yu. Olshanskii, SQ-*universality of hyperbolic groups* (Russian), Mat. Sb. **186** (1995), no. 8, 119–132; translation in Sb. Math. **186** (1995), no. 8, 1199–1211.

[64] A.Yu. Olshanskii, *On the Bass-Lubotzky question about quotients of hyperbolic groups*, J. Algebra **226** (2000), no. 2, 807–817.

[65] A.Yu. Olshanskii, *On residualing homomorphisms and G–subgroups of hyperbolic groups*, Int. J. Alg. Comp., **3** (1993), 4, 365–409.

[66] A.Yu. Olshanskii, M.V. Sapir, *The conjugacy problem for groups, and Higman embeddings*, Electron. Res. Announc. Amer. Math. Soc. **9** (2003), 40–50.

[67] D.V. Osin, *Weak hyperbolicity and free constructions*, Contemp. Math., to appear.

[68] D.V. Osin, *Relatrively hyperbolic groups and embedding theorems,* preprint, 2004.

[69] D.V. Osin, M.V. Sapir, *Asymptotic cones of relatively hyperbolic groups*, preprint, 2002.

[70] A. Pankrat'ev, *Hyperbolic products of groups,* Vestnik Mosk. Universiteta, Ser. 1, Mathematics, Mechanics, **2** (1999), 9–13 (in Russian).

[71] A. Pankrat'ev, *On the infinite hyperbolic quotients of hyperbolic products of groups*, Fundam. Prikl. Mat. 7 (2001), no. 2, 465–493.

[72] A. Pankrat'ev, *Hyperbolic products of groups,* PhD Thesises, Moscow State University, 2001.

[73] P. Papasoglou, *On the sub–quadratic isoperimetric inequality,* Geometric Group Theory, (R. Charney, M. Davis, M. Shapiro, ed.), de Gruyter, Berlin – New–York, 1995, 149–158.

[74] I. Rips, ...

[75] Z. Sela, *Endomorphisms of hyperbolic groups I: The Hopf property,* Topology, **38** (1999), 301–322.

[76] Z. Sela, *Diophantine geometry over groups I: Makanin – Razborov diagrams*, IHES Publ. Math., **93** (2001), 31–105.

[77] J-P. Serre, Trees, Springer–Verlag, 1980, Translation of "Arbres, Amalgames, SL_2", Astérisque, **46** 1977.

[78] H. Short, *Groups and combings*, preprint, ENS Lion, 1990.

[79] H. Short (editor), *Notes on word hyperbolic groups,* Group Theory from a Geometric Viewpoint, (E. Ghys, A. Haefliger, A. Verjovsky, ed.), Proc. ICTP Trieste 1990, World Scientific, Singapore, 1991, 3–64.

[80] A. Szczepański, *Relatively hyperbolic groups,* Michigan Math. J., **45** (1998), 611–618.

[81] P. Tukia, *Convergence groups and Gromov's metric hyperbolic spaces*, New Zeland J. Math, **23** (1994), 157–187.

[82] D. Wise, Non–positively curved squared complexes, aperiodic tillings, and non–residually finite groups, Ph.D. Thesis, Princeton Univ., 1996.

[83] D. Wise, *The residual finiteness of negatively curved polygons of finite groups*, Invent. Math. **149** (2002), no. 3, 579–617.

[84] A. Yaman, *A topplogical characterization of relatively hyperbolic groups*, prep., 2002.